

APPLICATION FOR UNITED STATES LETTERS PATENT
for
DYEING COMPOSITION COMPRISING A CATIONIC
PYRROLIDINE-BASED PARA-PHENYLENEDIAMINE, A TERTIARY
PARA-PHENYLENEDIAMINE AND A BENZOMORPHOLINE COUPLER,
METHODS AND USES

by
Francois COTTARD
and
Christine RONDEAU

<u>EXPRESS MAIL MAILING LABEL</u>	
NUMBER	EV 323285848 US
DATE OF DEPOSIT	December 12, 2003

CROSS REFERENCE TO RELATED APPLICATIONS

5 This application claims priority to French Application No. 02/15774 filed 13 December 2003, and further claims the benefit of U.S. Provisional Application No. 60/450,331 filed 28 February 2003, the entire disclosures of which are incorporated herein by reference in their entirety.

BACKGROUND OF THE INVENTION

10 The subject of the present application is a dyeing composition for dyeing keratinous fibres, in particular human keratinous fibres such as hair, comprising, in an appropriate dyeing medium, at least one cationic tertiary para-phenylenediamine containing a pyrrolidine ring, at least one non-cationic tertiary para-phenylenediamine, and at
15 least one benzomorpholine coupler.

The subject of the invention is also the use of this composition for dyeing keratinous fibres and the dyeing method using this composition.

20 It is known to dye keratinous fibres, and in particular human hair, with dyeing compositions containing oxidation dye precursors, generally called oxidation bases, such as ortho- or para-phenylenediamines, or ortho- or para-aminophenols and heterocyclic compounds. These oxidation bases are colourless or weakly coloured compounds which, when combined with oxidizing products, can give
25 rise, through a process of oxidative condensation, to coloured compounds.

It is also known that it is possible to vary the shades obtained with these oxidation bases by combining them with couplers or colour modifiers, the latter being chosen in particular from aromatic meta-
30 diamines, meta-aminophenols, meta-diphenols and certain heterocyclic compounds such as indole compounds.

The variety of molecules used in the oxidation bases and couplers allows a rich palette of colours to be obtained.

35 The so-called "permanent" coloration obtained using these oxidation dyes should moreover satisfy a number of requirements. Thus, it should be without disadvantage from the toxicological point of view, it should make it possible to obtain shades in the desired

intensity and should exhibit good fastness to external agents such as light, adverse weather conditions, washing, permanent waving, perspiration and rubbing.

5 The dyes should also make it possible to cover grey hair, and should finally be as less selective as possible, that is to say it is possible to obtain the least possible colour variations along the length of the same keratinous fibre, which is in general differentially sensitized (that is to say damaged) between its tip and its root.

10 It has already been proposed, in patent application WO 02/45675, to use compositions for the oxidation dyeing of keratinous fibres comprising a cationic tertiary para-phenylenediamine containing a pyrrolidine ring.

15 These cationic tertiary para-phenylenediamines containing a pyrrolidine ring lead to compositions which exhibit a harmlessness which is generally better than the compositions containing conventional para-phenylenediamines. However, the shades obtained when these compositions are used are markedly less intense and markedly more selective, that is to say that the dyes obtained exhibit substantial variations in colorations as a function of the degree of sensitization of the various types of hair or of the various areas of the same hair. The fastness of these shades can also vary greatly according to the degree of sensitization. In addition, when only tertiary para-phenylenediamines, including cationic tertiary para-phenylenediamines, are used, it is difficult to produce natural shades. 20 The shades obtained when such tertiary para-phenylenediamines are used are generally more ashen, that is to say less chromatic and poorly covering grey hair. 25

SUMMARY OF THE INVENTION

30 Surprisingly and advantageously, the applicant has just discovered that it is possible to obtain novel compositions for dyeing keratinous fibres, in particular human keratinous fibres such as hair, capable of overcoming the disadvantages above and, in particular, of leading to colorations with natural shades which are varied, chromatic, 35 aesthetic, not very selective and which withstand well the various attacks to which the fibres may be subjected, by combining, in the same composition, at least one cationic tertiary para-phenylenediamine

containing a pyrrolidine ring, at least one non-cationic tertiary para-phenylenediamine and at least one benzomorpholine coupler.

In addition, these compositions exhibit a good toxicological profile.

5 The subject of the invention is also a dyeing method using this composition, and a multicompartment dyeing device or dyeing kit.

Another subject of the invention is the use of the composition of the present invention for dyeing keratinous fibres, in particular human keratinous fibres such as hair.


10 The composition of the present invention makes it possible in particular to obtain a chromatic coloration of keratinous fibres which is very intense, little selective and fast while avoiding degradation of these fibres.

15 For the purposes of the present invention, cationic tertiary para-phenylenediamine containing a pyrrolidine ring is understood to mean a para-phenylenediamine possessing an NH₂ group and in the para position thereof a di-substituted amine functional group whose substitutions form with the nitrogen a pyrrolidine ring, the molecule possessing at least one quaternized nitrogen atom.

20 For the purposes of the present invention, non-cationic tertiary para-phenylenediamine is understood to mean a para-phenylenediamine possessing an NH₂ group and, at the para position thereof, a di-substituted amine functional group, the molecule not possessing a quaternized nitrogen atom.

25 In the context of the present invention, the expression alkyl is understood to mean linear or branched radicals, for example methyl, ethyl, n-propyl, isopropyl, butyl and the like. An alkoxy radical is an alk-O radical, the alkyl radical having the definition above. Halogen preferably denotes Cl, Br, I, F.

30 Among the cationic tertiary para-phenylenediamines containing a pyrrolidine ring which can be used in the composition according to the present invention, there may be mentioned in particular the compounds of the following formula (I) and their addition salts.



in which

- n varies from 0 to 4, it being understood that when n is greater than or equal to 2, then the radicals R_1 may be identical or different,
- 5 • R_1 represents a halogen atom; a saturated or unsaturated, aliphatic or alicyclic, C_1 - C_6 hydrocarbon chain, it being possible for the chain to contain one or more oxygen, nitrogen, silicon or sulphur atoms or an SO_2 group, and it being possible for the chain to be substituted with one or more hydroxyl or amino radicals; an onium radical Z , the radical R_1 not
- 10 containing a peroxide bond, or diazo, nitro or nitroso radicals,
- R_2 represents an onium radical Z or a radical $-X-C=NR_8-NR_9R_{10}$ in which X represents an oxygen atom or a radical $-NR_{11}$ and R_8 , R_9 , R_{10} and R_{11} represent a hydrogen atom, a C_1 - C_4 alkyl radical or a C_1 - C_4
- 15 hydroxyalkyl radical,
- R_3 represents a hydrogen atom or a hydroxyl radical.

Onium denotes the quaternary radical of a nitrogenous base.


By way of example, R_1 may be a chlorine atom, a methyl, ethyl, isopropyl, vinyl, allyl, methoxymethyl, hydroxyethyl, 1-carboxymethyl, 20 1-aminomethyl, 2-carboxyethyl, 2-hydroxyethyl, 3-hydroxypropyl, 1,2-dihydroxyethyl, 1-hydroxy-2-aminoethyl, 1-amino-2-hydroxyethyl, 1,2-diaminoethyl, methoxy, ethoxy, allyloxy, or 2-hydroxyethyloxy radical.

In particular, n is equal to 0.

In formula (I), when n is equal to 1, R_1 is preferably a halogen atom; a saturated or unsaturated, aliphatic or alicyclic, C_1 - C_6 hydrocarbon chain, it being possible for one or more carbon atoms to be replaced with an oxygen, nitrogen, silicon or sulphur atom, or with an SO_2 group, the radical R_1 not containing a peroxide bond, or diazo, nitro or nitroso radicals. Preferably, R_1 is chosen from chlorine,

bromine, C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl, C₁-C₄ aminoalkyl, C₁-C₄ alkoxy and C₁-C₄ hydroxyalkoxy radicals. By way of example, R₁ is chosen from a methyl, hydroxymethyl, 2-hydroxyethyl, 1,2-dihydroxyethyl, methoxy, isopropoxy or 2-hydroxyethoxy radical.

The radical R₂ of formula (I) is, according to a particular embodiment, the onium radical Z corresponding to formula (II)



in which

- D is a single bond of a linear or branched C₁-C₁₄ alkylene chain which may contain one or more heteroatoms chosen from oxygen, sulphur or nitrogen, and which may be substituted with one or more hydroxyl, C₁-C₆ alkoxy or amino radicals and which may carry one or more ketone functional groups;

- R₄, R₅ and R₆, taken separately, represent a C₁-C₁₅ alkyl radical; a C₁-C₆ monohydroxyalkyl radical; a C₂-C₆ polyhydroxyalkyl radical; a (C₁-C₆)alkoxy(C₁-C₆)alkyl radical; an aryl radical; a benzyl radical; a C₁-C₆ amidoalkyl radical; a tri(C₁-C₆)alkylsilane(C₁-C₆)alkyl radical; a C₁-C₆ aminoalkyl radical; a C₁-C₆ aminoalkyl radical in which the amine is mono- or di-substituted with a C₁-C₄ alkyl, (C₁-C₆)alkylcarbonyl, amido or (C₁-C₆)alkylsulphonyl radical; or

- R₄, R₅ and R₆ together, in pairs, form, with the nitrogen atom to which they are attached, a 4-, 5-, 6- or 7-membered saturated carbon ring which may contain one or more heteroatoms such as, for example, an azetidine ring, a pyrrolidine ring, a piperidine ring, a piperazine ring, or a morpholine ring, it being possible for the cationic ring to be substituted with a halogen atom, a hydroxyl radical, a C₁-C₆ alkyl radical, a C₁-C₆ monohydroxyalkyl radical, a C₂-C₆ polyhydroxyalkyl radical, a C₁-C₆ alkoxy radical, a tri(C₁-C₆)alkylsilane(C₁-C₆)alkyl

radical, an amido radical, a carboxyl radical, a (C₁-C₆)alkylcarbonyl radical, a thio (-SH) radical, a C₁-C₆ thioalkyl (-R-SH) radical, a (C₁-C₆)alkylthio radical, an amino radical, an amino radical which is mono- or di-substituted with a (C₁-C₆)alkyl, (C₁-C₆)alkylcarbonyl, amido or (C₁-C₆)alkylsulphonyl radical;

• R₇ represents a C₁-C₆ alkyl radical; a C₁-C₆ monohydroxyalkyl radical; a C₂-C₆ polyhydroxyalkyl radical; an aryl radical; a benzyl radical; a C₁-C₆ aminoalkyl radical; a C₁-C₆ aminoalkyl radical whose amine is mono- or di-substituted with a (C₁-C₆)alkyl, (C₁-C₆)alkylcarbonyl, amido or (C₁-C₆)alkylsulphonyl radical; a C₁-C₆ carboxyalkyl radical; a C₁-C₆ carbamylalkyl radical; a C₁-C₆ trifluoroalkyl radical; a tri(C₁-C₆)alkylsilane(C₁-C₆)alkyl radical; a C₁-C₆ sulphonamidoalkyl radical; a (C₁-C₆)alkylcarboxy(C₁-C₆)alkyl radical; a (C₁-C₆)alkylsulphinyl(C₁-C₆)alkyl radical; a (C₁-C₆)alkylsulphonyl(C₁-C₆)alkyl radical; a (C₁-C₆)alkylcarbonyl(C₁-C₆)alkyl radical; an N-(C₁-C₆)alkylcarbamyl(C₁-C₆)alkyl radical; an N-(C₁-C₆)alkylsulphonamido(C₁-C₆)alkyl radical;

• x is 0 or 1,
- when x = 0, then the linking arm is attached to the nitrogen atom carrying the radicals R₄ to R₆;

when x = 1, then two of the radicals R₄ to R₆ form, together with the nitrogen atom to which they are attached, a 4-, 5-, 6- or 7-membered saturated ring and D is linked to the carbon atom of the saturated ring;

• Y is a counter-ion.

In formula (II), when x is equal to 0, then R₄, R₅ and R₆ separately are preferably chosen from a C₁-C₆ alkyl radical, a C₁-C₄ monohydroxyalkyl radical, a C₂-C₄ polyhydroxyalkyl radical, a (C₁-C₆)alkoxy(C₁-C₄)alkyl radical, a C₁-C₆ amidoalkyl radical, a tri(C₁-C₆)alkylsilane(C₁-C₆)alkyl radical, or R₄ with R₅ form together an azetidine ring, a pyrrolidine, piperidine, piperazine or morpholine ring, R₆ being chosen in this case from a C₁-C₆ alkyl radical; a C₁-C₆ monohydroxyalkyl radical, a C₂-C₆ polyhydroxyalkyl radical; a C₁-C₆ aminoalkyl radical, an aminoalkyl radical which is mono- or di-substituted with a (C₁-C₆)alkyl radical, a (C₁-C₆)alkylcarbonyl, amido or (C₁-C₆)alkylsulphonyl radical; a C₁-C₆ carbamylalkyl radical; a tri(C₁-C₆)alkylsilane(C₁-C₆)alkyl radical; a (C₁-C₆)alkyl carboxy(C₁-


C₆)alkyl radical; a (C₁-C₆)alkylcarbonyl(C₁-C₆)alkyl radical; an N-(C₁-C₆)alkylcarbamyl(C₁-C₆)alkyl radical.

When x is equal to 1, then R₇ is preferably chosen from a C₁-C₆ alkyl radical; a C₁-C₆ monohydroxyalkyl radical; a C₂-C₆ polyhydroxy alkyl radical, a C₁-C₆ aminoalkyl radical, a C₁-C₆ aminoalkyl radical whose amine is mono- or di-substituted with a (C₁-C₆)alkyl, (C₁-C₆)alkylcarbonyl, amido or (C₁-C₆)alkylsulphonyl radical; a C₁-C₆ carbamylalkyl radical, a tri(C₁-C₆)alkylsilane(C₁-C₆)alkyl radical; a (C₁-C₆)alkylcarboxy(C₁-C₆)alkyl radical; a (C₁-C₆)alkylcarbonyl(C₁-C₆)alkyl radical; an N-(C₁-C₆)alkylcarbamyl(C₁-C₆)alkyl radical; R₄ with R₅ together form an azetidine, pyrrolidine, piperidine, piperazine or morpholine ring, R₆ being chosen in this case from a C₁-C₆ alkyl radical; a C₁-C₆ monohydroxyalkyl radical; a C₂-C₆ polyhydroxyalkyl radical; a C₁-C₆ aminoalkyl radical, a C₁-C₆ aminoalkyl radical whose amine is mono- or di-substituted with a (C₁-C₆)alkyl, (C₁-C₆)alkylcarbonyl, amido or (C₁-C₆)alkylsulphonyl radical; a C₁-C₆ carbamylalkyl radical; a tri(C₁-C₆)alkylsilane(C₁-C₆)alkyl radical; a (C₁-C₆)alkylcarboxy(C₁-C₆)alkyl radical; a (C₁-C₆)alkylcarbonyl(C₁-C₆)alkyl radical; an N-(C₁-C₆)alkylcarbamyl(C₁-C₆)alkyl radical.

In the formula (II), D is preferably a single bond or an alkylene chain which may be substituted.

When the radical R₂ corresponds to formula (II), it is preferably a trialkylammonium radical whose alkyl radicals may be substituted.

According to a second embodiment, the radical R₂ represents the onium radical Z corresponding to formula (III)



(III)

in which

D is a single bond or a linear or branched C₁-C₁₄ alkylene chain which may contain one or more heteroatoms chosen from oxygen, sulphur or nitrogen, and which may be substituted with one or more hydroxyl, C₁-C₆ alkoxy or amino radicals, and which may carry one or more ketone functional groups;

the vertices E, G, J, L, which are identical or different, represent a carbon, oxygen, sulphur or nitrogen atom to form a pyrrole, pyrazole, imidazole, triazole, oxazole, isooxazole, thiazole, isothiazole ring,

q is an integer between 0 and 4 inclusive;

o is an integer between 0 and 3 inclusive;

q+o is an integer between 0 and 4;

the radicals R₈, which are identical or different, represent a halogen atom, a hydroxyl radical, a C₁-C₆ alkyl radical, a C₁-C₆ monohydroxyalkyl radical, a C₂-C₆ polyhydroxyalkyl radical, a C₁-C₆ alkoxy radical, a tri(C₁-C₆)alkylsilane(C₁-C₆)alkyl radical, an amido radical, a carboxyl radical, a C₁-C₆ alkylcarbonyl radical, a thio radical, a C₁-C₆ thioalkyl radical, a (C₁-C₆)alkylthio radical, an amino radical, an amino radical which is mono- or di-substituted with a (C₁-C₆)alkyl, (C₁-C₆)alkylcarbonyl, amido or (C₁-C₆)alkylsulphonyl radical; a C₁-C₆ monohydroxyalkyl radical or a C₂-C₆ polyhydroxyalkyl radical; it being understood that the radicals R₈ are carried by a carbon atom,

• the radicals R₉, which are identical or different, represent a C₁-C₆ alkyl radical, a C₁-C₆ monohydroxyalkyl radical, a C₂-C₆ polyhydroxyalkyl radical, a tri(C₁-C₆)alkylsilane(C₁-C₆)alkyl radical, a (C₁-C₆)alkoxy(C₁-C₆)alkyl radical, a C₁-C₆ carbamylalkyl radical, a (C₁-C₆)alkylcarboxy(C₁-C₆)alkyl radical, a benzyl radical; it being understood that the radicals R₉ are carried by a nitrogen,

• R₁₀ represents a C₁-C₆ alkyl radical; a C₁-C₆ monohydroxyalkyl radical; a C₂-C₆ polyhydroxyalkyl radical; an aryl radical; a benzyl radical; a C₁-C₆ aminoalkyl radical, a C₁-C₆ aminoalkyl radical whose amine is substituted with a (C₁-C₆)alkyl, (C₁-C₆)alkylcarbonyl, amido or (C₁-C₆)alkylsulphonyl radical; a C₁-C₆ carboxyalkyl radical; a C₁-C₆ carbamylalkyl radical; a C₁-C₆ trifluoroalkyl radical; a tri(C₁-C₆)alkylsilane(C₁-C₆)alkyl radical; a C₁-C₆ sulphonamidoalkyl radical; a (C₁-C₆)alkylcarboxy(C₁-C₆)alkyl radical; a (C₁-C₆)alkylsulphinyl(C₁-C₆)alkyl radical; a (C₁-C₆)alkylsulphonyl(C₁-C₆)alkyl radical; a (C₁-C₆)alkylsulphonyl(C₁-C₆)alkyl radical;

C₆)alkylcarbonyl(C₁-C₆)alkyl radical; an N-(C₁-C₆)alkylcarbamyl(C₁-C₆)alkyl radical; an N-(C₁-C₆)alkylsulphonamido(C₁-C₆)alkyl radical;

- x is 0 or 1

when x = 0, the linking arm D is attached to the nitrogen atom,


5 when x = 1, the linking arm D is attached to one of the vertices E, G, J or L,

- Y is a counter-ion.

The vertices E, G, J and L preferably form an imidazole ring.

10 Among the radicals R₂ of formulae (III), the preferred radicals are those in which x is equal to 0, D is a single bond or an alkylene chain which may be substituted.

According to a third embodiment, R₂ represents the onium radical Z corresponding to formula (IV)



(IV)

in which:

15 • D is a single bond or a linear or branched C₁-C₁₄ alkylene chain which may contain one or more heteroatoms chosen from an oxygen, sulphur or nitrogen atom, and which may be substituted with one or more hydroxyl, C₁-C₆ alkoxy or amino radicals, and which may carry one or more ketone functional groups;

20 • the vertices E, G, J, L and M, which are identical or different, represent a carbon, oxygen, sulphur or nitrogen atom to form a ring chosen from the pyridine, pyrimidine, pyrazine, triazine and pyridazine rings;

- p is an integer between 0 and 3 inclusive;
- m is an integer between 0 and 5 inclusive;
- p+m is an integer between 0 and 5;

• the radicals R_{11} , which are identical or different, represent a halogen atom, a hydroxyl radical, a C_1 - C_6 alkyl radical, a C_1 - C_6 monohydroxyalkyl radical, a C_2 - C_6 polyhydroxyalkyl radical, a C_1 - C_6 alkoxy radical, a tri(C_1 - C_6)alkylsilane(C_1 - C_6)alkyl radical, an amido radical, a carboxyl radical, a C_1 - C_6 alkylcarbonyl radical, a thio radical, a C_1 - C_6 thioalkyl radical, a (C_1 - C_6)alkylthio radical, an amino radical, an amino radical which is substituted with a (C_1 - C_6)alkyl, (C_1 - C_6)alkylcarbonyl, amido or (C_1 - C_6)alkylsulphonyl radical; a C_1 - C_6 monohydroxyalkyl radical or a C_2 - C_6 polyhydroxyalkyl radical; it being understood that the radicals R_{11} are carried by a carbon atom,

• the radicals R_{12} , which are identical or different, represent a C_1 - C_6 alkyl radical, a C_1 - C_6 monohydroxyalkyl radical, a C_2 - C_6 polyhydroxyalkyl radical, a tri(C_1 - C_6)alkylsilane(C_1 - C_6)alkyl radical, a (C_1 - C_6)alkoxy(C_1 - C_6)alkyl radical, a C_1 - C_6 carbamylalkyl radical, a (C_1 - C_6)alkylcarboxy(C_1 - C_6)alkyl radical, a benzyl radical; it being understood that the radicals R_{12} are carried by a nitrogen,

• R_{13} represents a C_1 - C_6 alkyl radical; a C_1 - C_6 monohydroxyalkyl radical; a C_2 - C_6 polyhydroxyalkyl radical; an aryl radical; a benzyl radical; a C_1 - C_6 aminoalkyl radical, a C_1 - C_6 aminoalkyl radical whose amine is mono- or di-substituted with a (C_1 - C_6)alkyl, (C_1 - C_6)alkylcarbonyl, amido or (C_1 - C_6)alkylsulphonyl radical; a C_1 - C_6 carboxyalkyl radical; a C_1 - C_6 carbamylalkyl radical; a C_1 - C_6 trifluoroalkyl radical; a tri(C_1 - C_6)alkylsilane(C_1 - C_6)alkyl radical; a C_1 - C_6 sulphonamidoalkyl radical; a (C_1 - C_6)alkylcarboxy(C_1 - C_6)alkyl radical; a (C_1 - C_6)alkylsulphinyl(C_1 - C_6)alkyl radical; a (C_1 - C_6)alkylsulphonyl(C_1 - C_6)alkyl radical; a (C_1 - C_6)alkylcarbonyl(C_1 - C_6)alkyl radical; an N-(C_1 - C_6)alkylcarbamyl(C_1 - C_6)alkyl radical; an N-(C_1 - C_6)alkylsulphonamido(C_1 - C_6)alkyl radical;

• x is 0 or 1
when $x = 0$, the linking arm D is attached to the nitrogen atom,
when $x = 1$, the linking arm D is attached to one of the vertices E, G, J, L or M,

• Y is a counter-ion.

Preferably, the vertices E, G, J, L and M form, with the nitrogen of the ring, a pyridine and pyrimidine ring

When x is equal to 0, then R_{11} is preferably chosen from a hydroxyl radical, a C_1 - C_6 alkyl radical, a C_1 - C_6 monohydroxyalkyl radical, a C_2 - C_6 polyhydroxyalkyl radical, a C_1 - C_6 alkoxy radical, a

tri(C₁-C₆)alkylsilane(C₁-C₆)alkyl radical, an amido radical, a C₁-C₆ alkylcarbonyl radical, an amino radical, an amino radical which is mono- or di-substituted with a (C₁-C₆)alkyl, a (C₁-C₆)alkylcarbonyl, amido or (C₁-C₆)alkylsulphonyl radical; a C₁-C₆ monohydroxyalkyl radical or a C₂-C₆ polyhydroxyalkyl radical and R₁₂ is chosen from a C₁-C₆ alkyl radical, a C₁-C₆ monohydroxyalkyl radical, a C₂-C₆ polyhydroxyalkyl radical, a tri(C₁-C₆)alkylsilane(C₁-C₆)alkyl radical, a (C₁-C₆)alkoxy(C₁-C₆)alkyl radical, a C₁-C₆ carbamylalkyl radical.

When x is equal to 1, R₁₃ is preferably chosen from a C₁-C₆ alkyl radical; a C₁-C₆ monohydroxyalkyl radical; a C₂-C₆ polyhydroxyalkyl radical; a C₁-C₆ aminoalkyl radical, a C₁-C₆ aminoalkyl radical whose amine is mono- or di-substituted with a (C₁-C₆)alkyl radical, a (C₁-C₆)alkylcarbonyl radical, an amido radical, a (C₁-C₆)alkylsulphonyl radical; a C₁-C₆ carbamylalkyl radical; a tri(C₁-C₆)alkylsilane(C₁-C₆)alkyl radical; a (C₁-C₆)alkylcarbonyl(C₁-C₆)alkyl radical; an N-(C₁-C₆)alkylcarbamyl(C₁-C₆)alkyl radical; R₁₁ is chosen from a hydroxyl radical, a C₁-C₆ alkyl radical, a C₁-C₆ monohydroxyalkyl radical, a C₂-C₆ polyhydroxyalkyl radical, a C₁-C₆ alkoxy radical, a tri(C₁-C₆)alkylsilane(C₁-C₆)alkyl radical, an amido radical, a C₁-C₆ alkylcarbonyl radical, an amino radical, an amino radical which is mono- or di-substituted with a (C₁-C₆)alkyl, (C₁-C₆)alkylcarbonyl, amido or (C₁-C₆)alkylsulphonyl radical; and R₁₂ is chosen from a C₁-C₆ alkyl radical, a C₁-C₆ monohydroxyalkyl radical, a C₂-C₆ polyhydroxyalkyl radical, a tri(C₁-C₆)alkylsilane(C₁-C₆)alkyl radical, a (C₁-C₆)alkoxy(C₁-C₆)alkyl radical, a C₁-C₆ carbamylalkyl radical.

Preferably, R₁₁, R₁₂ and R₁₃ are alkyl radicals which may be substituted.

The radical R₂ may also represent an onium radical of formula
-XP(O)(O-)OCH₂CH₂N⁺(CH₃)₃
where X represents an oxygen atom or a radical -NR₁₄, R₁₄ representing a hydrogen, a C₁-C₄ alkyl radical or a hydroxyalkyl radical.

In the context of the invention, R₂ may also represent a guanidine radical of formula -X-C=NR₈-NR₉R₁₀, X represents an oxygen atom or a radical -NR₁₁, R₈, R₉, R₁₀ and R₁₁ representing a hydrogen, a C₁-C₄ alkyl radical or a hydroxyalkyl radical. According









to a particular embodiment, X is $-NR_{11}$, R_8 is a hydrogen, R_9 and R_{10} are chosen from hydrogen or an alkyl, preferably methyl, radical.







The pKa of the guanidine radical R_2 is in general such that this substituent is present in cationic form ($=NR_8H^+$) under conventional conditions for oxidation hair dyeing.









In the context of the invention, the counter-ion may be derived from a halogen atom such as bromine, chlorine, fluorine or iodine, a hydroxide, a citrate, a succinate, a tartrate, a lactate, a tosylate, a mesylate, a benzenesulphonate, an acetate, a hydrogen sulphate or a C_1 - C_6 alkyl sulphate such as for example methyl sulphate or ethyl sulphate.









In the context of the present application, cationic tertiary para-phenylenediamines containing a pyrrolidine ring, which are described above and for which R_2 is of formula II or III, are preferably used. Still more preferably, the cationic tertiary para-phenylenediamines containing a pyrrolidine ring, which are described above and for which R_2 is of formula II or of formula III, with $x=0$ and for which $n=0$, are used.









By way of example of derivatives of formula (I), there may be mentioned:









Formula	Nomenclature	Formula	Nomenclature
	[1-(4-Aminophenyl)pyrrolidin-3-yl]-trimethylammonium chloride (1)		[1-(4-Aminophenyl)pyrrolidin-3-yl]dimethyltetradecylammonium bromide (2)
	N'-[1-(4-Aminophenyl)pyrrolidin-3-yl]-N,N-dimethylguanidinium chloride (3)		N-[1-(4-Aminophenyl)pyrrolidin-3-yl]guanidinium chloride (4)
	3-[1-(4-Aminophenyl)pyrrolidin-3-yl]-1-methyl-3H-imidazole-1-ium chloride (5)		[1-(4-Aminophenyl)pyrrolidin-3-yl]-(2-hydroxyethyl)dimethylammonium chloride (6)
	[1-(4-Aminophenyl)pyrrolidin-3-yl]dimethyl-(3-trimethylsilylpropyl)ammonium chloride		[1-(4-Aminophenyl)pyrrolidin-3-yl]-(trimethylammoniumhexyl)dimethylammonium dichloride (8)







	(7)		
	[1-(4-Aminophenyl)-pyrrolidin-3-yl]oxophosphoryl choline (9)		{2-[1-(4-Aminophenyl)-pyrrolidin-3-yloxy]-ethyl}-trimethylammonium chloride (10)
	1-{2-[1-(4-Aminophenyl)pyrrolidin-3-yloxy]ethyl}-1-methylpyrrolidinium; chloride (11)		3-{3-[1-(4-Aminophenyl)pyrrolidin-3-yloxy]-propyl}-1-methyl-3H-imidazol-1-ium; chloride (12)
	1-{2-[1-(4-Aminophenyl)pyrrolidin-3-yloxy]ethyl}-1-methylpiperidinium; chloride (13)		3-{3-[1-(5-trimethylsilyl-4-amino-3-trimethylsilyl-ethylphenyl)pyrrolidin-3-yloxy]propyl}-1-methyl-3H-imidazole-1-um; chloride (14)

	<p>[1-(4-Amino-3-methylphenyl)pyrrolidin-3-yl]-trimethylammonium chloride (15)</p>		<p>[1-(4-Amino-3-methylphenyl)pyrrolidin-3-yl]dimethyl-tetradecylammonium chloride (16)</p>
	<p>N'-[1-(4-Amino-3-methylphenyl)-pyrrolidin-3-yl]-N,N-dimethylguanidinium chloride (17)</p>		<p>N-[1-(4-Amino-3-methylphenyl)-pyrrolidin-3-yl]-guanidinium chloride (18)</p>
	<p>3-[1-(4-Amino-3-methylphenyl)-pyrrolidin-3-yl]-1-methyl-3H-imidazole-1-ium chloride (19)</p>		<p>[1-(4-Amino-3-methylphenyl)-pyrrolidin-3-yl]-(2-hydroxyethyl)-dimethylammonium chloride (20)</p>
	<p>[1-(4-Amino-3-methylphenyl)pyrrolidin-3-yl]-dimethyl(3-trimethylsilylpropyl)ammonium chloride (21)</p>		<p>[1-(4-Amino-3-methylphenyl)pyrrolidin-3-yl]-(trimethylammonium-hexyl)dimethylammonium dichloride (22)</p>

	[1-(4-Amino-3-methylphenyl)-pyrrolidin-3-yl]-oxophosphorylcholine (23)		{2-[1-(4-Amino-3-methylphenyl)-pyrrolidin-3-yloxy]ethyl}-trimethylammonium chloride (24)
	1-{2-[1-(4-Amino-3-methylphenyl)pyrrolidin-3-yloxy]ethyl}-1-methylpyrrolidinium chloride (25)		3-{3-[1-(4-Amino-3-methylphenyl)-pyrrolidin-3-yloxy]propyl}-1-methyl-3H-imidazole-1-ium chloride (26)
	1-{2-[1-(4-Amino-3-methylphenyl)pyrrolidin-3-yloxy]ethyl}-1-methylpiperidinium chloride (27)		[1-(4-Amino-3-trimethylsilylphenyl)-pyrrolidin-3-yl]-trimethylammonium chloride (28)
	3-[1-(4-Amino-3-trimethylsilylphenyl)pyrrolidin-3-yl]-1-methyl-3H-imidazole-1-ium chloride (29)		3-{3-[1-(4-Amino-3-trimethylsilylphenyl)-pyrrolidin-3-yloxy]propyl}-1-methyl-3H-imidazole-1-ium chloride (30)

	<p>[1-(5-trimethylsilyl-4-amino-3-trimethylsilylphenyl)pyrrolidin-3-yl]-trimethylammoniumchloride (31)</p>		<p>3-[1-(5-trimethylsilyl-4-amino-3-trimethylsilylphenyl)-pyrrolidin-3-yl]-1-methyl-3H-imidazole-1-ium chloride (32)</p>
	<p>1'-(4-Aminophenyl)-1-methyl-[1,3']bipyrrolidinyl-1-ium chloride (33)</p>		<p>1'-(4-Amino-3-methylphenyl)-1-methyl-[1,3']bipyrrolidinyl-1-ium chloride (34)</p>
	<p>3-[[1-(4-Aminophenyl)pyrrolidin-3-yl]carbonyl]-1-methyl-3H-imidazole-1-ium chloride (35)</p>		<p>3-[[1-(4-Amino-3-methylphenyl)pyrrolidin-3-yl]carbonyl]-1-methyl-3H-imidazole-1-ium chloride (36)</p>
	<p>3-[1-(4-Aminophenyl)pyrrolidin-3-yl]-1-(3-trimethylsilylpropyl)-3H-imidazole-1-ium chloride</p>		<p>3-[1-(4-Aminophenyl)pyrrolidin-3-yl]-1-(3-trimethylsilylpropyl)-3H-imidazole-1-ium chloride (38)</p>

	(37)		
	[1-(4-Aminophenyl)pyrrolidin-3-yl]-ethyl-dimethylammonium chloride (39)		[1-(4-Aminophenyl)pyrrolidin-3-yl]-ethyl-dimethylammonium iodide (40)
	[1-(4-aminophenyl)pyrrolidin-3-yl]-propyl-dimethylammonium iodide, (41)		[1-(4-Aminophenyl)pyrrolidin-3-yl]-propyl-dimethylammonium bromide (42)
	[1-(4-Aminophenyl)pyrrolidin-3-yl]-propyl-dimethylammonium methosulphate (43)		[1-(4-Aminophenyl)pyrrolidin-3-yl]-butyl-dimethylammonium iodide (44)
	[1-(4-Aminophenyl)pyrrolidin-3-yl]-pentyl-dimethylammonium iodide (45)		[1-(4-Aminophenyl)pyrrolidin-3-yl]-hexyl-dimethylammonium iodide (46)

	[1-(4-Aminophenyl)pyrrolidin-3-yl]-heptyldimethylammonium iodide (47)		[1-(4-Aminophenyl)pyrrolidin-3-yl]-octyldimethylammonium iodide (48)
	[1-(4-Aminophenyl)pyrrolidin-3-yl]-decyldimethylammonium iodide (49)		[1-(4-aminophenyl)pyrrolidin-3-yl]-hexadecyldimethylammonium iodide (50)
	[1-(4-Aminophenyl)pyrrolidin-3-yl]-hydroxyethyl-dimethylammonium chloride (51)		[1-(4-aminophenyl)-pyrrolidin-3-yl]-hydroxyethyl-dimethylammonium iodide (52)

The derivatives of formula I which are preferably used are:

[1-(4-Aminophenyl)pyrrolidin-3-yl]trimethylammonium chloride;

5 [1-(4-Aminophenyl)pyrrolidin-3-yl]dimethyltetradecylammonium bromide;

N'-[1-(4-Aminophenyl)pyrrolidin-3-yl]-N,N-dimethylguanidinium choride

10 N-[1-(4-Aminophenyl)pyrrolidin-3-yl] guanidinium choride
3-[1-(4-Aminophenyl)pyrrolidin-3-yl]-1-methyl-3H-imidazol-1-ium chloride;

[1-(4-Aminophenyl)pyrrolidin-3-yl]-(2-hydroxyethyl)dimethylammonium chloride

[1-(4-Aminophenyl)pyrrolidin-3-yl]dimethyl-(3-trimethylsilanylpropyl)ammonium chloride;

[1-(4-Amino-3-methylphenyl)pyrrolidin-3-yl]trimethylammonium chloride

5 [1-(4-Amino-3-methylphenyl)pyrrolidin-3-yl]dimethyl-tetradecylammonium chloride

N'-[1-(4-Amino-3-methylphenyl)pyrrolidin-3-yl]-N,N-dimethylguanidinium choride

10 N-[1-(4-Amino-3-methylphenyl)pyrrolidin-3-yl] guanidinium choride

3-[1-(4-Amino-3-methylphenyl)pyrrolidin-3-yl]-1-methyl-3H-imidazol-1-ium chloride

[1-(4-Amino-3-methylphenyl)pyrrolidin-3-yl]-(2-hydroxyethyl)dimethylammonium chloride

15 [1-(4-Amino-3-methylphenyl)pyrrolidin-3-yl]dimethyl-(3-trimethylsilanylpropyl)ammonium chloride

1'-(4-Aminophenyl)-1-methyl-[1,3']bipyrrolidinyl-1-ium chloride

1'-(4-Amino-3-methylphenyl)-1-methyl-[1,3']bipyrrolidinyl-1-ium chloride

20 3-{[1-(4-Aminophenyl)pyrrolidin-3-ylcarbamoyl]methyl}-1-methyl-3H-imidazol-1-ium chloride

3-{[1-(4-Amino-3-methylphenyl)pyrrolidin-3-ylcarbamoyl]methyl}-1-methyl-3H-imidazol-1-ium chloride

25 3-[1-(4-Aminophenyl)pyrrolidin-3-yl]-1-(3-trimethylsilanylpropyl)-3H-imidazol-1-ium chloride

3-[1-(4-Amino-3-methylphenyl)pyrrolidin-3-yl]-1-(3-trimethylsilanylpropyl)-3H-imidazol-1-ium chloride

[1-(4-aminophenyl)pyrrolidin-3-yl]ethyldimethylammonium chloride

30 [1-(4-aminophenyl)pyrrolidin-3-yl]-ethyldimethylammonium iodide

[1-(4-aminophenyl)pyrrolidin-3-yl]propyldimethylammonium iodide,

35 [1-(4-aminophenyl)pyrrolidin-3-yl]propyldimethylammonium bromide

[1-(4-aminophenyl)pyrrolidin-3-yl]propyldimethylammonium methosulphate

[1-(4-aminophenyl)pyrrolidin-3-yl]butyldimethylammonium
 iodide
 [1-(4-aminophenyl)pyrrolidin-3-yl]pentyldimethylammonium
 iodide
 5 [1-(4-aminophenyl)pyrrolidin-3-yl]hexyldimethylammonium
 iodide
 [1-(4-aminophenyl)pyrrolidin-3-yl]heptyldimethylammonium
 iodide
 [1-(4-aminophenyl)pyrrolidin-3-yl]octyldimethylammonium
 10 iodide
 [1-(4-aminophenyl)pyrrolidin-3-yl]decyldimethylammonium
 iodide
 [1-(4-aminophenyl)pyrrolidin-3-yl]hexadecyldimethyl-
 ammonium iodide
 15 [1-(4-aminophenyl)pyrrolidin-3-yl]hydroxyethyldimethyl-
 ammonium chloride
 [1-(4-aminophenyl)pyrrolidin-3-yl]hydroxyethyldimethyl-
 ammonium iodide.
 More preferably, the following compounds will be used:
 20 [1-(4-Aminophenyl)pyrrolidin-3-yl]trimethylammonium
 chloride
 [1-(4-Aminophenyl)pyrrolidin-3-yl]dimethyltetradecyl-
 ammonium bromide
 N'-[1-(4-Aminophenyl)pyrrolidin-3-yl]-N,N-dimethyl
 25 guanidinium choride
 N-[1-(4-Aminophenyl)pyrrolidin-3-yl] guanidinium choride
 3-[1-(4-Aminophenyl)pyrrolidin-3-yl]-1-methyl-3H-imidazol-1-
 ium chloride
 [1-(4-Aminophenyl)pyrrolidin-3-yl]-(2-
 30 hydroxyethyl)dimethylammonium chloride
 [1-(4-Aminophenyl)pyrrolidin-3-yl]dimethyl-(3-
 trimethylsilanylpropyl)ammonium chloride
 [1-(4-Aminophenyl)pyrrolidin-3-yl]-(trimethylammonium-
 hexyl)dimethylammonium dichloride
 35 1'-(4-Aminophenyl)-1-methyl[1,3']bipyrrolidinyl-1-ium
 chloride
 3-[1-(4-Aminophenyl)pyrrolidin-3-yl]-1-(3-trimethylsilanyl-
 propyl)-3H-imidazol-1-ium chloride

3-[1-(4-Amino-3-methylphenyl)pyrrolidin-3-yl]-1-(3-trimethylsilanylpropyl)-3H-imidazol-1-ium chloride

[1-(4-aminophenyl)pyrrolidin-3-yl]ethyldimethylammonium chloride

5 [1-(4-aminophenyl)pyrrolidin-3-yl]ethyldimethylammonium iodide

[1-(4-aminophenyl)pyrrolidin-3-yl]propyldimethylammonium iodide,

[1-(4-aminophenyl)pyrrolidin-3-yl]propyldimethylammonium bromide

10 [1-(4-aminophenyl)pyrrolidin-3-yl]propyldimethylammonium methosulphate

[1-(4-aminophenyl)pyrrolidin-3-yl]butyldimethylammonium iodide

15 [1-(4-aminophenyl)pyrrolidin-3-yl]pentyldimethylammonium iodide

[1-(4-aminophenyl)pyrrolidin-3-yl]hexyldimethylammonium iodide

[1-(4-aminophenyl)pyrrolidin-3-yl]heptyldimethylammonium iodide

20 [1-(4-aminophenyl)pyrrolidin-3-yl]octyldimethylammonium iodide

[1-(4-aminophenyl)pyrrolidin-3-yl]decyldimethylammonium iodide

25 [1-(4-aminophenyl)pyrrolidin-3-yl]hexadecyldimethylammonium iodide

[1-(4-aminophenyl)pyrrolidin-3-yl]hydroxyethyldimethylammonium chloride

[1-(4-aminophenyl)pyrrolidin-3-yl]hydroxyethyldimethylammonium iodide

30 Still more preferably, the following compounds will be used:

[1-(4-Aminophenyl)pyrrolidin-3-yl]trimethylammonium chloride

35 3-[1-(4-Aminophenyl)pyrrolidin-3-yl]-1-methyl-3H-imidazol-1-ium chloride

[1-(4-Aminophenyl)pyrrolidin-3-yl]-(2-hydroxyethyl)-dimethylammonium chloride

1'-(4-Aminophenyl)-1-methyl[1,3']bipyrrolidinyl-1-ium
chloride, and in particular

[1-(4-Aminophenyl)pyrrolidin-3-yl]trimethylammonium
chloride, and


5 [1-(4-Aminophenyl)pyrrolidin-3-yl]-(2-hydroxyethyl)-
dimethylammonium chloride.

10 The counter-ion is not critical as for the result of the invention,
any compounds similar to the preferred compounds described above
but with a different counter-ion forms an integral part of the preferred
compounds.

The cationic tertiary para-phenylenediamine(s) containing a
pyrrolidine ring represent from 0.001% to 10%, and preferably from
0.005% to 6% by weight relative to the total weight of the
composition.

15 The compounds of formula (I) may be synthesized according to
known methods, and in particular methods described in application
WO 02/45675.

20 The non-cationic tertiary para-phenylenediamines which can be
used for the purposes of the present application preferably correspond
to the compounds of general formula (V) and their addition salts with
an acid:



25 in which R₁ is a linear or branched C₁-C₆ alkyl radical or a
linear or branched C₂-C₆ hydroxyalkyl radical,

R₂ is a linear or branched C₂-C₆ hydroxyalkyl radical,

R₃ denotes a hydrogen atom or a linear or branched C₁-C₆
alkyl radical, a C₁-C₆ alkoxy radical or a halogen atom, and

30 R₄ is a radical situated at any of the free positions of the
benzene ring and denotes a hydrogen atom or a linear or branched
C₁-C₆ alkyl radical, a linear or branched C₁-C₆ alkoxy radical or a
halogen atom.

These compounds and their synthesis are described in application DE 2 240 495.

When R'₁, R'₃ or R'₄ denote a linear or branched C₁-C₆ alkyl radical, it is preferably a radical chosen from methyl, ethyl, n-propyl, isopropyl, n-butyl, s-butyl, tert-butyl n-amyl, isoamyl and n-hexyl radicals.

When R'₁ or R'₂ denote a linear or branched C₂-C₆ hydroxyalkyl radical, it is preferably a radical chosen from monohydroxyalkyl, dihydroxyalkyl and trihydroxyalkyl radicals. More particularly, they are 2-hydroxyethyl, 3-hydroxypropyl, 2-hydroxypropyl, tris(hydroxymethyl)methyl, 1,3-dihydroxy-2-methylpropyl, 2,3-dihydroxypropyl and 1,3-dihydroxy-2-propyl radicals.


When R'₃ and R'₄ denote a halogen, it is Cl, Br, I, or F.

When R'₃ or R'₄ denote a linear or branched C₁-C₆ alkoxy radical, it is preferably a radical chosen from methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, n-amyloxy, isoamyloxy and n-hexyloxy radicals.

Preferably, the non-cationic tertiary para-phenylenediamine used in the composition according to the present application is N,N-bis(β-hydroxyethyl)-p-phenylenediamine whose synthesis is described in patent US 3 884 627.

According to the present invention, the non-cationic tertiary para-phenylenediamines represent from 0.001 to 15%, and preferably from 0.05 to 10% by weight relative to the total weight of the composition.

The benzomorpholine couplers which can be used in the compositions according to the present application correspond to the compounds of formula (V') and their addition salts with an acid:



in which X represents a sulphur (S) atom or an oxygen (O) atom, R'' represents a hydroxyl or amino group at the 5, 6 or 7 position, R''₁ and R''₂ represent a hydrogen atom, a linear or branched

C₁-C₆ alkyl radical or a linear or branched C₂-C₆ hydroxyalkyl radical, it being possible for the benzene ring to additionally contain another linear or branched C₁-C₆ alkyl radical.

5 When R''₁ and R''₂ denote a linear or branched C₁-C₆ alkyl radical, it is preferably a radical chosen from methyl, ethyl, n-propyl, isopropyl, n-butyl, s-butyl, tert-butyl, n-amyl, isoamyl and n-hexyl radicals.

10 When R''₁ or R''₂ denote a linear or branched C₂-C₆ hydroxyalkyl radical, it is preferably a radical chosen from monohydroxyalkyl, dihydroxyalkyl and trihydroxyalkyl radicals. More particularly, they are 2-hydroxyethyl, 3-hydroxypropyl, 2-hydroxypropyl, tris(hydroxymethyl)methyl, 1,3-dihydroxy-2-methylpropyl, 2,3-dihydroxypropyl and 1,3-dihydroxy-2-propyl radicals.


Preferably, X denotes an oxygen atom.

15 More particularly, the following compounds are used as benzomorpholine couplers: 7-hydroxybenzomorpholine, 7-aminobenzomorpholine, 6-hydroxybenzomorpholine, 6-aminobenzomorpholine, 5-hydroxybenzomorpholine, 5-aminobenzomorpholine, 1-methyl-7-hydroxybenzomorpholine, 2-methyl-7-hydroxybenzomorpholine, 5-hydroxy-7-methylbenzomorpholine. Preferably, 20 6-hydroxybenzomorpholine is used.

According to the present invention, the benzomorpholine couplers represent from 0.001 to 15%, and preferably from 0.05 to 10% by weight relative to the total weight of the composition.

25 Advantageously, the compositions according to the present application comprise at least one cationic tertiary para-phenylenediamine containing a pyrrolidine ring, at least one non-cationic tertiary para-phenylenediamine, at least one benzomorpholine coupler and at least one dye chosen from the para-aminophenols of formula V'' and the heterocyclic couplers of formula (V''') and 30 mixtures thereof.

The para-aminophenols which can be used in the compositions according to the present application are the compounds of formula (V'') and their addition salts



5 in which:

- R'₅ represents a hydrogen or halogen atom, a C₁-C₄ alkyl, C₁-C₄ monohydroxyalkyl, (C₁-C₄)alkoxy(C₁-C₄)alkyl, C₁-C₄ aminoalkyl or hydroxy(C₁-C₄)alkylamino(C₁-C₄)alkyl radical,
- R'₆ represents a hydrogen or halogen atom, a C₁-C₄ alkyl, C₁-C₄ monohydroxyalkyl, C₂-C₄ polyhydroxyalkyl, C₁-C₄ aminoalkyl, C₁-C₄ cyanoalkyl or (C₁-C₄)alkoxyl(C₁-C₄)alkyl radical,
- R'_a represents a hydrogen atom or a C₁-C₄ alkyl radical.


10 When R'₅ or R'₆ denotes a C₁-C₄ alkyl radical, it is preferably a radical chosen from methyl, ethyl, n-propyl, isopropyl, n-butyl, s-butyl and tert-butyl radicals.

The expression (C₁-C₄) "alkoxy" is understood to mean a (C₁-C₄)alkyl-O radical, the alkyl radical having the definition given above.

When R'₅ or R'₆ denotes a halogen, it is Cl, Br, I or F.

20 More particularly, the compounds of formula (V'') are chosen from para-aminophenol, 4-amino-3-methylphenol, 4-amino-3-fluorophenol, 4-amino-3-hydroxymethylphenol, 4-amino-2-methylphenol, 4-amino-2-hydroxymethylphenol, 4-amino-2-methoxymethylphenol, 4-amino-2-aminomethylphenol, 4-amino-2-(β-hydroxyethylamino-methyl)phenol, 4-amino-2-fluorophenol, N-methyl-para-aminophenol and their addition salts.

The heterocyclic couplers which can be used in the compositions according to the present application are the compounds of formula (V''') and their addition salts.



in which OH occupies positions 6 or 7 of the aromatic ring and R'₇ denotes a hydrogen atom, a C₁-C₄ alkyl radical; R'₈ and R'₉, which are identical or different, denote a hydrogen atom, a lower C₁-C₄ alkyl radical, a carboxyl radical or a (C₁-C₄)alkoxycarbonyl radical, and their salts.

Among the compounds of formula V''', the compounds which are particularly preferred are the compounds in which the alkyl radical denotes methyl or ethyl; the alkoxycarbonyl radical denotes methoxy or ethoxycarbonyl.

Among these compounds, there may be mentioned 6-hydroxyindole, 6-hydroxy-3-methoxycarbonylindole, 6-hydroxy-1-methyl-3-methoxycarbonylindole, 6-hydroxy-1-methyl-2,3-dimethoxycarbonylindole, 6-hydroxy-1,2-dimethylindole, 6-hydroxy-2-methylindole, 6-hydroxy-2-carboxyindole, 6-hydroxy-2,3-dimethylindole, 6-hydroxy-3-carboxyindole, 6-hydroxy-3-ethoxycarbonylindole, 6-hydroxy-2-ethoxycarbonylindole, 6-hydroxy-3-methylindole, 6-hydroxy-1-methylindole, 7-hydroxyindole, 7-hydroxy-3-methylindole.

Preferably, 6-hydroxyindole is used.

According to the present application, the compounds of formulae (V'') and/or (V''') represent from 0.0001 to 10%, and preferably 0.001 to 5% by weight relative to the total weight of the composition.

In general, the addition salts of the compounds of formulae (V), (V'), (V'') and (V''') are chosen from hydrochlorides, hydrobromides, sulphates, citrates, succinates, tartrates, lactates, tosylates, benzenesulphonates, phosphates and acetates.

According to a first preferred embodiment, the composition according to the present invention additionally contains at least one cationic polymer.

For the purposes of the present invention, the expression "cationic polymer" denotes any polymer containing cationic groups and/or groups which are ionizable to cationic groups.

The cationic polymers which can be used in accordance with the present invention may be chosen from all those already known per se as improving the cosmetic properties of hair, namely in particular

those described in patent application EP-A-337 354 and in French patents FR-2 270 846, 2 383 660, 2 598 611, 2 470 596 and 2 519 863.





The preferred cationic polymers are chosen from those which contain units comprising primary, secondary, tertiary and/or quaternary amine groups which may either be part of the main polymer chain, or which may be carried by a side substituent directly linked to the latter.

The cationic polymers used generally have a number-average molecular mass between 500 and $5 \cdot 10^6$ approximately, and preferably between 10^3 and $3 \cdot 10^6$ approximately.

Among the cationic polymers, there may be mentioned more particularly polymers of the polyamine, polyamino amide and poly(quaternaryammonium) type.

They are known products. They are described in particular in French patents No. 2 505 348 or 2 542 997. Among the said polymers, there may be mentioned:

the homopolymers or copolymers derived from acrylic or methacrylic esters or amides and comprising at least one of the units of the following formulae (VI), (VII), (VIII) or (IX):



in which:

R₃ denotes a hydrogen atom or a CH₃ radical;

A represents a linear or branched alkyl group of 1 to 6 carbon atoms, preferably 2 or 3 carbon atoms or a hydroxyalkyl group of 1 to 4 carbon atoms;

5 R₄, R₅, R₆, which are identical or different, represent an alkyl group having from 1 to 18 carbon atoms or a benzyl radical and preferably an alkyl group having from 1 to 6 carbon atoms;

R₁ and R₂, which are identical or different, represent hydrogen or an alkyl group having from 1 to 6 carbon atoms and preferably methyl or ethyl;

10 X⁻ denotes an anion derived from an inorganic or organic acid such as a methosulphate anion or a halide such as chloride or bromide.

The polymers of the family (1) may contain, in addition, one or more units derived from comonomers which may be chosen from the family of acrylamides, methacrylamides, diacetone acrylamides, acrylamides and methacrylamides substituted on the nitrogen with lower (C₁-C₄)alkyls, acrylic or methacrylic acids or esters thereof, vinyl lactams such as vinylpyrrolidone or vinylcaprolactam, vinyl esters.

20 Thus, among these polymers of the family (1), there may be mentioned:

- the copolymers of acrylamide and dimethylaminoethyl methacrylate quaternized with dimethyl sulphate or with a dimethyl halide such as that sold under the name HERCOFLOC by the company HERCULES,

- the copolymers of acrylamide and methacryloyloxyethyltrimethylammonium chloride described, for example, in Patent Application EP-A-080976 and sold under the name BINA QUAT P 100 by the company CIBA GEIGY,

- the copolymer of acrylamide and methacryloyloxyethyltrimethylammonium methosulphate sold under the name RETEN by the company HERCULES,

- the vinylpyrrolidone/dialkylaminoalkyl acrylate or methacrylate copolymers, quaternized or otherwise, such as the products sold under the name "GAFQUAT" by the company ISP such as for example "GAFQUAT 734" or "GAFQUAT 755" or alternatively

the products called "COPOLYMER 845, 958 and 937". These polymers are described in detail in French Patents 2 077 143 and 2 393 573,

5 - the dimethylaminoethyl methacrylate/vinylcaprolactam/vinylpyrrolidone terpolymers such as the product sold under the name GAFFIX VC 713 by the company ISP,

 - the vinylpyrrolidone/methacrylamidopropyl dimethylamine copolymers marketed in particular under the name STYLEZE CC 10 by ISP,

10 - and the quaternized vinylpyrrolidone/dimethyl aminopropyl methacrylamide copolymers such as the product sold under the name "GAFQUAT HS 100" by the company ISP.

 (2) The cellulose ether derivatives comprising quaternary ammonium groups, described in French Patent 1 492 597, and in particular the polymers marketed under the names "JR" (JR 400, JR 125, JR 30M) or "LR" (LR 400, LR 30M) by the company Union Carbide Corporation. These polymers are also defined in the CTFA dictionary as hydroxyethyl cellulose quaternary ammoniums which have reacted with an epoxide substituted by a trimethylammonium group.

20 (3) Cationic cellulose derivatives such as cellulose copolymers or cellulose derivatives grafted with a quaternary ammonium water-soluble monomer, and described especially in US Patent 4 131 576, such as hydroxyalkyl celluloses like hydroxymethyl, hydroxyethyl or hydroxypropyl celluloses grafted especially with a

25 methacryloylethyltrimethylammonium, methacrylamidopropyl-trimethylammonium or dimethyldiallylammonium salt.

 The commercialized products corresponding to this definition are more particularly the products sold under the name "Celquat L 200" and "Celquat H 100" by the company National Starch.

30 (4) The cationic polysaccharides described more particularly in US Patents 3 589 578 and 4 031 307 such as guar gums containing cationic trialkylammonium groups. Guar gums modified with a 2,3-epoxypropyltri methylammonium salt (e.g. chloride) are for example used.

35 Such products are marketed in particular under the trade names JAGUAR C13 S, JAGUAR C 15, JAGUAR C 17 or JAGUAR C162 by the company MEYHALL.

(5) Polymers consisting of piperazinyl units and of alkylene or hydroxyalkylene divalent radicals with straight or branched chains, optionally interrupted by oxygen, sulphur or nitrogen atoms or by aromatic or heterocyclic rings, as well as the oxidation and/or quaternization products of these polymers. Such polymers are described especially in French patents 2 162 025 and 2 280 361;

(6) Water-soluble polyaminoamides prepared in particular by polycondensation of an acid compound with a polyamine; these polyaminoamides may be crosslinked with an epihalohydrin, a diepoxide, a dianhydride, an unsaturated dianhydride, a diunsaturated derivative, a bishalohydrin, a bisazetidinium, a bishaloacyldiamine, an alkylbishalide or else with an oligomer resulting from the reaction of a difunctional compound which is reactive towards a bishalohydrin, a bisazetidinium, a bishaloacyldiamine, an alkylbishalide, an epihalohydrin, a diepoxide or a diunsaturated derivative; the crosslinking agent being employed in proportions ranging from 0.025 to 0.35 mol per amine group of the polyaminoamide; these polyaminoamides may be alkylated or, if they include one or more tertiary amine functional groups, quaternized. Such polymers are described especially in French Patents 2 252 840 and 2 368 508.

(7) Polyaminoamide derivatives resulting from the condensation of polyalkylenepolyamines with polycarboxylic acids, followed by an alkylation with difunctional agents. There may be mentioned, for example, the adipic acid/dialkylaminohydroxy-alkyldialkylenetriamine polymers in which the alkyl radical contains from 1 to 4 carbon atoms and preferably denotes methyl, ethyl or propyl. Such polymers are described especially in French Patent 1 583 363.



Among these derivatives there may be mentioned more particularly the adipic acid/dimethyl aminohydroxy-propyl/diethylenetriamine polymers sold under the name "Cartaretine F, F4 or F8" by the company Sandoz.

(8) Polymers obtained by reaction of a polyalkylenepolyamine containing two primary amine groups and at least one secondary amine group with a dicarboxylic acid chosen from diglycolic acid and saturated aliphatic dicarboxylic acids containing from 3 to 8 carbon atoms. The molar ratio of the polyalkylenepolyamine to the dicarboxylic acid being between 0.8 : 1 and 1.4 : 1; the

polyaminoamide resulting therefrom being made to react with epichlorohydrin in a molar ratio of epichlorohydrin relative to the secondary amine group of the polyaminoamide of between 0.5 : 1 and 1.8 : 1. Such polymers are described especially in American Patents 3 227 615 and 2 961 347.

Polymers of this type are marketed in particular under the name "Hercosett 57" by the company Hercules Inc. or else under the name of "PD 170" or "Delsette 101" by the company Hercules in the case of the copolymer of adipic acid/epoxypropyl/diethylene triamine.

(9) Cyclopolymers of alkyldiallylamine or of dialkyldiallylammmonium, such as the homopolymers or copolymers comprising, as main constituent of the chain, units corresponding to the formulae (X) or (XI):




in which formulae k and t are equal to 0 or 1, the sum k + t being equal to 1; R₉ denotes a hydrogen atom or a methyl radical; R₇ and R₈, independently of each other, denote an alkyl group containing from 1 to 6 carbon atoms, a hydroxyalkyl group in which the alkyl group preferably has 1 to 5 carbon atoms, or a lower (C₁-C₄)amidoalkyl group or R₇ and R₈ may denote, jointly with the nitrogen atom to which they are attached, heterocyclic groups such as piperidinyl or morpholinyl; R₇ and R₈, independently of each other, preferably denote an alkyl group having 1 to 4 carbon atoms; Y⁻ is an

anion such as bromide, chloride, acetate, borate, citrate, tartrate, bisulphate, bisulphite, sulphate or phosphate. These polymers are described especially in French Patent 2 080 759 and in its certificate of addition 2 190 406.

5 Among the polymers defined above there may be mentioned more particularly the dimethyldiallylammonium chloride homopolymer sold under the name "Merquat 100" by the company Calgon (and its homologues of low weight-average molecular masses) and the copolymers of diallyl dimethylammonium chloride and acrylamide
10 marketed under the name "MERQUAT 550".

(10) The quaternary diammonium polymer containing repeat units corresponding to the formula:



15 formula (XII) in which:

R₁₀, R₁₁, R₁₂ and R₁₃, which are identical or different, represent aliphatic, alicyclic or arylaliphatic radicals containing from 1 to 6 carbon atoms or lower hydroxyalkyl aliphatic radicals, or else
20 R₁₀, R₁₁, R₁₂ and R₁₃, together or separately, form, with the nitrogen atoms to which they are attached, heterocyclic rings optionally containing a second heteroatom other than nitrogen, or else R₁₀, R₁₁, R₁₂ and R₁₃ denote a linear or branched C₁-C₆ alkyl radical substituted by a nitrile, ester, acyl, amide or -CO-O-R₁₄-D or -CO-NH-R₁₄-D
25 group where R₁₄ is an alkylene and D a quaternary ammonium group;

A₁ and B₁ represent polymethylene groups containing from 2 to 20 carbon atoms which may be linear or branched, saturated or unsaturated and which may contain, bonded to or inserted into the main chain, one or more aromatic rings, or one or more oxygen or
30 sulphur atoms or sulfoxide, sulphone, disulphide, amino, alkylamino, hydroxyl, quaternary ammonium, ureido, amide or ester groups, and

X⁻ denotes an anion derived from an inorganic or organic acid;
A₁, R₁₀ and R₁₂, with the two nitrogen atoms to which they are attached, may form a piperazine ring; in addition if A₁ denotes a
35 saturated or unsaturated, linear or branched alkylene or

hydroxyalkylene radical, B₁ may also denote a group -(CH₂)_n-CO-D-OC-(CH₂)_n- in which D denotes:

a) a glycol residue of formula: -O-Z-O-, where Z denotes a linear or branched hydrocarbon radical or a group corresponding to one of the following formulae:



where x and y denote an integer from 1 to 4, representing a defined and unique degree of polymerization or any number from 1 to 4 representing a mean degree of polymerization;

b) a disecundary diamine residue such as a piperazine derivative;

c) a diprimary diamine residue of formula: -NH-Y-NH-, where Y denotes a linear or branched hydrocarbon radical or else the divalent radical




d) a ureylene group of formula: -NH-CO-NH-;

X- is preferably an anion such as chloride or bromide.

These polymers have a number-average molecular mass which is generally between 1 000 and 100 000.


Polymers of this type are described especially in French Patents 2 320 330, 2 270 846, 2 316 271, 2 336 434 and 2 413 907 and US Patents 2 273 780, 2 375 853, 2 388 614, 2 454 547, 3 206 462, 2 261 002, 2 271 378, 3 874 870, 4 001 432, 3 929 990, 3 966 904, 4 005 193, 4 025 617, 4 025 627, 4 025 653, 4 026 945 and 4 027 020.

It is possible to use more particularly the polymers which consist of repeat units corresponding to the following formula (XIII):



in which R₁₀, R₁₁, R₁₂ and R₁₃, which are identical or different, denote an alkyl or hydroxyalkyl radical having from 1 to 4 carbon atoms approximately, n and p are integers varying from 2 to 20 approximately and X- is an anion derived from an inorganic or organic acid.

(11) The quaternary polyammonium polymers consisting of repeat units of formula XIV):



(XIV)

5 in which p denotes an integer varying from 1 to 6 approximately, D may be zero or may represent a group $-(\text{CH}_2)_r\text{---CO-}$ in which r denotes a number equal to 4 or to 7, X⁻ is an anion.

Such polymers may be prepared according to the methods described in USA Patents No. 4 157 388, 4 702 906, 4 719 282. They
10 are in particular described in patent application EP-A-122 324.

Among these, there may be mentioned for example the products "Mirapol A 15", "Mirapol AD1", "Mirapol AZ1" and Mirapol 175" sold by the company Miranol.

(12) Quaternary vinylpyrrolidone and vinylimidazole polymers
15 such as, for example, the products marketed under the names Luviquat FC 905, FC 550 and FC 370 by the company B.A.S.F.


(13) Polyamines like the Polyquart H sold by HENKEL, referenced under the name of "Polyethylene glycol (15) Tallow Polyamine" in the CTFA dictionary.

20 (14) The crosslinked polymers of methacryloyloxy(C₁-C₄ alkyl)tri(C₁-C₄ alkyl)ammonium salts such as the polymers obtained by homopolymerization of dimethylaminoethyl methacrylate quaternized with methyl chloride, or by copolymerization of acrylamide with dimethylaminoethyl methacrylate quaternized with
25 methyl chloride, the homo- or copolymerization being followed by crosslinking with a compound containing olefinic unsaturation, in particular methylenebisacrylamide. More particularly, it is possible to employ a crosslinked acrylamide/methacryloyloxyethyltrimethylammonium chloride copolymer (20/80 by weight) in the form of a
30 dispersion containing 50% by weight of the said copolymer in mineral oil. This dispersion is marketed under the name of "SALCARE® SC


92" by the company ALLIED COLLOIDS. It is also possible to employ a crosslinked methacryloyloxyethyltrimethylammonium chloride homopolymer containing approximately 50% by weight of the homopolymer in mineral oil or in a liquid ester. These dispersions are marketed under the names of "SALCARE® SC 95" and "SALCARE® SC 96" by the company ALLIED COLLOIDS.

Other cationic polymers that may be employed within the scope of the invention are polyalkyleneimines, in particular polyethyleneimines, polymers containing vinylpyridine or vinylpyridinium units, condensates of polyamines and of epichlorohydrin, quaternary polyureylenes and chitin derivatives.

Among all the cationic polymers which may be used in the context of the present invention, it is preferable to use the polymers of the families (1), (9), (10), (11) and (14) and more preferably still the polymers with the repeat units of the following formulae (W) and (U):



and in particular those whose molecular weight, determined by gel permeation chromatography, is between 9 500 and 9 900;



and in particular those whose molecular weight, determined by gel permeation chromatography, is about 1 200.

The cationic polymer concentration in the composition according to the present invention may vary from 0.01 to 10% by weight relative to the total weight of the composition, preferably from 0.05 to 5% and more preferably still from 0.1 to 3%.

According to a second preferred embodiment, the composition according to the present invention additionally contains at least one thickening polymer also called "rheology-adjusting agents".

5 The rheology-adjusting agents may be chosen from fatty acid amides (diethanol- or monoethanolamide of copra, monoethanolamide of oxyethylenated alkyl ether carboxylic acid), cellulosic thickeners (hydroxyethylcellulose, hydroxypropylcellulose, carboxymethyl-cellulose), guar gum and its derivatives (hydroxypropylguar), gums of microbial origin (xanthan gum, scleroglucan gum), crosslinked
10 homopolymers of acrylic acid or of acrylamidopropanesulphonic acid and the associative polymers as described below.

The associative polymers which can be used according to the invention are water-soluble polymers which are capable, in an aqueous medium, of reversibly combining with each other or with other
15 molecules.


Their chemical structure comprises hydrophilic regions, and hydrophobic regions which are characterized by at least one fatty chain.

The associative polymers which can be used according to the invention may be of the anionic, cationic, amphoteric and preferably nonionic type.
20

Their concentration by weight in the dyeing composition may vary from about 0.01 to 10% of the total weight of the composition and in the ready-to-use composition (comprising the oxidizing agent)
25 from about 0.0025 to 10% of the total weight of the composition. More preferably, this quantity varies from about 0.1 to 5% by weight in the dyeing composition and from about 0.025 to 10% in the ready-to-use composition.

Among the associative polymers of the anionic type, there may
30 be mentioned:

-(I) those comprising at least one hydrophilic unit and at least one allyl ether unit containing a fatty chain, more particularly those in which the hydrophilic unit consists of an ethylenic unsaturated anionic monomer, more particularly still of a vinylcarboxylic acid and most
35 particularly of an acrylic acid, a methacrylic acid or mixtures thereof, and in which the allyl ether unit containing a fatty chain corresponds to the monomer having the following formula (XV):



in which R' denotes H or CH₃, B denotes the ethyleneoxy radical, n is zero or denotes an integer ranging from 1 to 100, R denotes a hydrocarbon radical chosen from the alkyl, arylalkyl, aryl, alkylaryl or cycloalkyl radicals, comprising from 8 to 30 carbon atoms, preferably 10 to 24, and more particularly still from 12 to 18 carbon atoms. A more particularly preferred unit of formula (XV) is a unit in which R' denotes H, n is equal to 10, and R denotes a stearyl (C₁₈) radical.


Anionic associative polymers of this type are described and prepared, according to a method of polymerization in emulsion, in patent EP-0 216 479.

Among these anionic associative polymers, the polymers formed from 20 to 60% by weight of acrylic acid and/or of methacrylic acid, from 5 to 60% by weight of lower alkyl (meth)acrylates, from 2 to 50% by weight of allyl ether containing a fatty chain of formula (XV), and from 0 to 1% by weight of a crosslinking agent which is a well-known copolymerizable polyethylenic unsaturated monomer such as diallyl phthalate, allyl (meth)acrylate, divinylbenzene, (poly)ethylene glycol dimethacrylate and methylenebisacrylamide, are particularly preferred according to the invention.


Among the latter, the crosslinked terpolymers of methacrylic acid, ethyl acrylate, polyethylene glycol (10 EO) stearyl alcohol ether (Steareth 10), in particular those sold by the company ALLIED COLLOIDS under the names SALCARE SC80® and SALCARE SC90® which are aqueous emulsions containing 30% of a crosslinked terpolymer of methacrylic acid, of ethyl acrylate and of steareth-10-allyl ether (40/50/10) are most particularly preferred.

-(II) those comprising at least one hydrophilic unit of the olefinic unsaturated carboxylic acid type and at least one hydrophobic unit of the (C₁₀-C₃₀)alkyl ester of unsaturated carboxylic acid type.

Preferably, these polymers are chosen from those whose hydrophilic unit of the olefinic unsaturated carboxylic acid type corresponds to the monomer having the following formula (XVI):



in which R₁ denotes H or CH₃ or C₂H₅, that is to say acrylic acid, methacrylic acid or ethacrylic acid units, and whose hydrophobic unit of the (C₁₀-C₃₀)alkyl ester of unsaturated carboxylic acid type corresponds to the monomer having the following formula (XVII):



in which R₂ denotes H or CH₃ or C₂H₅ (that is to say acrylate, methacrylate or ethacrylate units) and preferably H (acrylate units) or CH₃ (methacrylate units), R₃ denoting a C₁₀-C₃₀, and preferably C₁₂-C₂₂, alkyl radical.

(C₁₀-C₃₀)alkyl esters of unsaturated carboxylic acids in accordance with the invention comprise for example lauryl acrylate, stearyl acrylate, decyl acrylate, isodecyl acrylate, dodecyl acrylate, and the corresponding methacrylates, lauryl methacrylate, stearyl methacrylate, decyl methacrylate, isodecyl methacrylate and dodecyl methacrylate.

Anionic polymers of this type are for example described and prepared according to patents US-3 915 921 and 4 509 949.

Among this type of anionic associative polymers, there will be more particularly used polymers formed from a mixture of monomers comprising:

(i) essentially acrylic acid,
(ii) an ester having the formula (XVII) described above and in which R₂ denotes H or CH₃, R₃ denoting an alkyl radical having from 12 to 22 carbon atoms,

(iii) and a crosslinking agent, which is a well-known copolymerizable polyethylenic unsaturated monomer such as diallyl phthalate, allyl (meth)acrylate, divinylbenzene, (poly)ethylene glycol dimethacrylate and methylenebisacrylamide.

Among this type of anionic associative polymers, there will be more particularly used those consisting of 95 to 60% by weight of acrylic acid (hydrophilic unit), 4 to 40% by weight of C₁₀-C₃₀ alkyl acrylate (hydrophobic unit), and 0 to 6% by weight of crosslinking polymerizable monomer, or alternatively those consisting of 98 to 96% by weight of acrylic acid (hydrophilic unit), 1 to 4% by weight of C₁₀-

C₃₀ alkyl acrylate (hydrophobic unit) and 0.1 to 0.6% by weight of crosslinking polymerizable monomer such as those described above.

Among said above polymers, the products sold by the company GOODRICH under the trade names PEMULEN TR1®, PEMULEN TR2®, CARBOPOL 1382®, and still more preferably PEMULEN TR1®, and the product sold by the company S.E.P.P.I.C. under the name COATEX SX®, are most particularly preferred according to the present invention.

-(III) the terpolymers of maleic anhydride/C₃₀-C₃₈ α-olefin/alkyl maleate such as the product (maleic anhydride/C₃₀-C₃₈ α-olefin/isopropyl maleate copolymer) sold under the name PERFORMA V 1608® by the company NEWPHASE TECHNOLOGIES.

-(IV) the acrylic terpolymers comprising:

(a) about 20% to 70% by weight of a carboxylic acid with α,β-monoethylenic unsaturation,

(b) about 20 to 80% by weight of a nonsurfactant monomer with α,β-monoethylenic unsaturation different from (a),

(c) about 0.5 to 60% by weight of a nonionic monourethane which is the product of the reaction of a monohydric surfactant with a monoisocyanate with monoethylenic unsaturation,

such as those described in patent application EP-A-0 173 109 and more particularly that described in Example 3, namely a methacrylic acid/methyl acrylate/dimethyl metaisopropenyl benzyl isocyanate of ethoxylated (40 EO) behenyl alcohol terpolymer in 25% aqueous dispersion.


-(V) the copolymers comprising among their monomers a carboxylic acid with α,β-monoethylenic unsaturation and an ester of a carboxylic acid with α,β-monoethylenic unsaturation and an oxyalkylenated fatty alcohol.

Preferably, these compounds also comprise, as monomer, an ester of a carboxylic acid with α,β-monoethylenic unsaturation and a C₁-C₄ alcohol.

By way of example of this type of compound, there may be mentioned ACULYN 22® sold by the company ROHM and HAAS, which is an oxyalkylenated stearyl methacrylate/ethyl acrylate/methacrylic acid terpolymer.

Among the associative polymers of the cationic type, there may be mentioned:

-(I) the cationic associative polyurethanes the family of which has been described by the applicant in French patent application No. 0009609; it can be represented by the following general formula (XVIII):



in which:

R and R', which are identical or different, represent a hydrophobic group or a hydrogen atom;

10 X and X', which are identical or different, represent a group containing an amine functional group carrying or otherwise a hydrophobic group, or alternatively the group L";

L, L' and L", which are identical or different, represent a group derived from a diisocyanate;

15 P and P', which are identical or different, represent a group containing an amine functional group carrying or otherwise a hydrophobic group;

Y represents a hydrophilic group;

r is an integer between 1 and 100, preferably between 1 and 50 and in particular between 1 and 25;

20 n, m and p are each, independently of the others, between 0 and 1000;

the molecule containing at least one protonated or quaternized amine functional group and at least one hydrophobic group.

25 In a preferred embodiment of these polyurethanes, the only hydrophobic groups are the groups R and R' at the chain ends.

A preferred family of cationic associative polyurethanes is that corresponding to the formula (XVIII) described above and in which:

R and R' both represent independently a hydrophobic group,

X, X' each represent a group L",

30 n and p are between 1 and 1000, and

L, L', L", P, P', Y and m have the meaning indicated above.

Another preferred family of cationic associative polyurethanes is that corresponding to the formula (XVIII) above in which:

35 R and R' both represent independently a hydrophobic group, X, X' each represent a group L", n and p are equal to 0, and L, L', L", Y and m have the meaning indicated above.

The fact that n and p are equal to 0 means that these polymers do not contain units derived from a monomer containing an amine

functional group, incorporated into the polymer during polycondensation. The protonated amine functional groups of these polyurethanes result from the hydrolysis of isocyanate functional groups, in excess, at the chain end, followed by alkylation of the
5 primary amine functional groups formed by alkylating agents containing a hydrophobic group, that is to say compounds of the RQ or R'Q type, in which R and R' are as defined above and Q denotes a leaving group such as a halide, a sulphate and the like.

Yet another preferred family of cationic associative
10 polyurethanes is that corresponding to the formula (Ia) above in which:

R and R' both represent independently a hydrophobic group,

X and X' both represent independently a group containing a quaternary amine,

15 n and p are equal to zero, and

L, L', Y and m have the meaning indicated above.


The number-average molecular mass of the cationic associative polyurethanes is preferably between 400 and 500 000, in particular between 1 000 and 400 000, and ideally between 1 000 and 300 000.

20 The expression hydrophobic group is understood to mean a radical or polymer containing a saturated or unsaturated, linear or branched hydrocarbon chain which may contain one or more heteroatoms such as P, O, N, S or a radical containing a perfluorinated or silicone chain. When it denotes a hydrocarbon radical, the
25 hydrophobic group contains at least 10 carbon atoms, preferably from 10 to 30 carbon atoms, in particular from 12 to 30 carbon atoms, and more preferably from 18 to 30 carbon atoms.

Preferably, the hydrocarbon group is derived from a monofunctional compound.

30 By way of example, the hydrophobic group may be derived from a fatty alcohol such as stearyl alcohol, dodecyl alcohol, decyl alcohol. It may also denote a hydrocarbon polymer such as for example polybutadiene.

When X and/or X' denote a group containing a tertiary or
35 quaternary amine, X and/or X' may represent one of the following formulae:




in which:

5 R_2 represents a linear or branched alkylene radical having from 1 to 20 carbon atoms, containing or otherwise a saturated or unsaturated ring, or an arylene radical, it being possible for one or more of the carbon atoms to be replaced by a heteroatom chosen from N, S, O, P;

10 R_1 and R_3 , which are identical or different, denote a linear or branched C_1 - C_{30} alkyl or alkenyl radical, an aryl radical, it being possible for at least one of the carbon atoms to be replaced by a heteroatom chosen from N, S, O, P;

A^- is a physiologically acceptable counter-ion.

The groups L, L' and L'' represent a group of formula:




in which:

Z represents -O-, -S- or -NH-; and

20 R_4 represents a linear or branched alkylene radical having from 1 to 20 carbon atoms, containing or otherwise a saturated or unsaturated ring, an arylene radical, it being possible for one or more of the carbon atoms to be replaced by a heteroatom chosen from N, S, O and P.

25 The groups P and P', comprising an amine functional group, may represent at least one of the following formulae:



in which:

R₅ and R₇ have the same meanings as R₂ defined above;

5 R₆, R₈ and R₉ have the same meanings as R₁ and R₃ defined above;

R₁₀ represents a linear or branched alkylene group, which is optionally unsaturated and which may contain one or more heteroatoms chosen from N, O, S and P,

10 and A⁻ is a physiologically acceptable counter-ion.

As regards the meaning of Y, the expression hydrophilic group is understood to mean a polymeric or nonpolymeric water-soluble group.

15 By way of example, there may be mentioned, when polymers are not involved, ethylene glycol, diethylene glycol and propylene glycol.


20 In the case, in accordance with a preferred embodiment, of a hydrophilic polymer, there may be mentioned, by way of example, polyethers, sulphonated polyesters, sulphonated polyamides, or a mixture of these polymers. Preferably, the hydrophilic compound is a polyether and in particular a polyethylene oxide or a polypropylene oxide.

The cationic associative polyurethanes of formula (XVIII) which can be used according to the invention are formed from diisocyanates and from various compounds possessing functional groups containing a labile hydrogen. The functional groups containing a labile hydrogen may be alcohol functional groups, primary or secondary amine functional groups or thiol functional groups which give, after reaction with the diisocyanate functional groups, polyurethanes, polyureas and polythioureas, respectively. The term "polyurethanes" which can be used according to the present invention covers these three types of polymer, namely polyurethanes proper, polyureas and polythioureas and copolymers thereof.


A first type of compounds entering into the preparation of the polyurethane of formula (XVIII) is a compound containing at least one unit containing an amine functional group. This compound may be multifunctional, but preferably the compound is difunctional, that is to say that according to a preferred embodiment, this compound contains two labile hydrogen atoms carried for example by a hydroxyl, primary amine, secondary amine or thiol functional group. It is also possible to use a mixture of multifunctional and difunctional compounds in which the percentage of multifunctional compounds is low.

As indicated above, this compound may contain more than one unit containing an amine functional group. It is then a polymer carrying a repeat of the unit containing an amine functional group.

This type of compounds may be represented by one of the following formulae:



or



in which Z, P, P', n and p are as defined above.

By way of example of a compound containing an amine functional group, there may be mentioned N-methyldiethanolamine, N-tert-butyldiethanolamine, N-sulphoethyldiethanolamine.

The second compound entering into the preparation of the polyurethane of formula (XVIII) is a diisocyanate corresponding to the formula:



in which R_4 is defined above.

By way of example, there may be mentioned methylenediphenyl diisocyanate, methylenecyclohexane diisocyanate, isophorone diisocyanate, toluene diisocyanate, naphthalene diisocyanate, butane diisocyanate, hexane diisocyanate.

5 A third compound entering into the preparation of the polyurethane of formula (XVIII) is a hydrophobic compound intended to form the terminal hydrophobic groups of the polymer of formula (XVIII).

10 This compound consists of a hydrophobic group and a functional group containing a labile hydrogen, for example a hydroxyl, primary or secondary amine, or thiol functional group.

 By way of example, this compound may be a fatty alcohol, such as in particular stearyl alcohol, dodecyl alcohol, decyl alcohol. When this compound contains a polymeric chain, it may be for example
15 α -hydroxyl -hydrogenated polybutadiene.

 The hydrophobic group of the polyurethane of formula (XVIII) may also result from the quaternization reaction of the tertiary amine of the compound containing at least one tertiary amine unit. Thus, the hydrophobic group is introduced by the quaternizing agent. This
20 quaternizing agent is a compound of the RQ or R'Q type, in which R and R' are as defined above and Q denotes a leaving group such as a halide, a sulphate, and the like.

 The cationic associative polyurethane may additionally comprise a hydrophilic sequence. This sequence is provided by a
25 fourth type of compound entering into the preparation of the polymer. This compound may be multifunctional. It is preferably difunctional. It is also possible to have a mixture where the percentage of multifunctional compound is low.

30 The functional groups containing a labile hydrogen are alcohol, primary or secondary amine, or thiol functional groups. This compound may be a polymer terminated at the chain ends by one of these functional groups containing a labile hydrogen.

 By way of example, there may be mentioned, when polymers are not involved, ethylene glycol, diethylene glycol and propylene
35 glycol.

 In the case of a hydrophilic polymer, there may be mentioned, by way of example, polyethers, sulphonated polyesters, sulphonated polyamides, or a mixture of these polymers. Preferably, the

hydrophilic compound is a polyether and in particular a polyethylene oxide or a polypropylene oxide.

The hydrophilic group noted Y in formula (XVIII) is optional. Indeed, the units containing a quaternary or protonated amine functional group may suffice to provide the solubility or water-dispersibility necessary for this type of polymer in an aqueous solution.

Although the presence of a hydrophilic group Y is optional, cationic associative polyurethanes are nevertheless preferred which contain such a group.

-(II) the quaternized cellulose derivatives and the polyacrylates with noncyclic amine-containing side groups.

The quaternized cellulose derivatives are in particular,

- the quaternized celluloses modified by groups comprising at least one fatty chain, such as the alkyl, arylalkyl or alkylaryl groups comprising at least 8 carbon atoms, or mixtures thereof,

- the quaternized hydroxyethylcelluloses modified by groups comprising at least one fatty chain, such as the alkyl, arylalkyl or alkylaryl groups comprising at least 8 carbon atoms, or mixtures thereof.

The alkyl radicals carried by the above quaternized celluloses or hydroxyethylcelluloses preferably comprise from 8 to 30 carbon atoms. The aryl radicals preferably denote phenyl, benzyl, naphthyl or anthryl groups.

There may be mentioned as examples of quaternized alkylhydroxyethylcelluloses containing C₈-C₃₀ fatty chains the products QUATRISOFT LM 200®, QUATRISOFT LM-X 529-18-A®, QUATRISOFT LM-X 529-18B® (C₁₂ alkyl) and QUATRISOFT LM-X 529-8® (C₁₈ alkyl) marketed by the company AMERCHOL and the products CRODACEL QM®, CRODACEL QL® (C₁₂ alkyl) and CRODACEL QS® (C₁₈ alkyl) marketed by the company CRODA.



Amphoteric associative polymers

The amphoteric associative polymers are preferably chosen from those containing at least one noncyclic cationic unit. Still more particularly, the ones that are preferred are those prepared from or comprising 1 to 20 mol% of monomer containing a fatty chain, and preferably 1.5 to 15 mol% and still more particularly 1.5 to 6 mol%, relative to the total number of moles of monomers.

The preferred amphoteric associative polymers according to the invention comprise, or are prepared by copolymerizing:

1) at least one monomer of formula (XIX) or (XX):

A⁻



5

in which R₁ and R₂, which are identical or different, represent a hydrogen atom or a methyl radical, R₃, R₄ and R₅, which are identical or different, represent a linear or branched alkyl radical having from 1 to 30 carbon atoms,

10

Z represents an NH group or an oxygen atom,

n is an integer from 2 to 5,

A⁻ is an anion derived from an organic or inorganic acid, such as a methosulphate anion or a halide such as chloride or bromide;

15

2) at least one monomer of formula (XXI)



in which R₆ and R₇, which are identical or different, represent a hydrogen atom or a methyl radical;

and

20

3) at least one monomer of formula (XXII):



in which R₆ and R₇, which are identical or different, represent a hydrogen atom or a methyl radical, X denotes an oxygen or nitrogen atom and R₈ denotes a linear or branched alkyl radical having from 1 to 30 carbon atoms;

25

at least one of the monomers of formula (XIX), (XX) or (XXII) containing at least one fatty chain.

The monomers of formula (XIX) and (XX) of the present invention are preferably chosen from the group consisting of:

30

- dimethylaminoethyl methacrylate, dimethylaminoethyl acrylate,
- diethylaminoethyl methacrylate, diethylaminoethyl acrylate,
- dimethylaminopropyl methacrylate, dimethylaminopropyl acrylate,

- dimethylaminopropylmethacrylamide,
dimethylaminopropylacrylamide,

these monomers being optionally quaternized, for example with
a C₁-C₄ alkyl halide or a C₁-C₄ dialkyl sulphate.

5 More particularly, the monomer of formula (XIX) is chosen
from acrylamidopropyltrimethylammonium chloride and
methacrylamidopropyltrimethylammonium chloride.

The monomers of formula (XXI) of the present invention are
preferably chosen from the group consisting of acrylic acid,
10 methacrylic acid, crotonic acid and 2-methylcrotonic acid. More
particularly, the monomer of formula (XXI) is acrylic acid.

The monomers of formula (XXII) of the present invention are
preferably chosen from the group consisting of C₁₂-C₂₂, and more
particularly C₁₆-C₁₈ alkyl acrylates or methacrylates.

15 The monomers constituting the amphoteric polymers containing
a fatty chain of the invention are preferably already neutralized and/or
quaternized.

The ratio of the number of cationic charges/anionic charges is
preferably equal to about 1.

20 The amphoteric associative polymers according to the
invention preferably comprise from 1 to 10 mol% of the monomer
containing a fatty chain (monomer of formula (XIX), (XX) or (XXII)),
and preferably from 1.5 to 6 mol%.

The weight-average molecular weights of the amphoteric
25 associative polymers according to the invention may vary from 500 to
50 000 000 and are preferably between 10 000 and 5 000 000.

The amphoteric associative polymers according to the
invention may also contain other monomers such as nonionic
monomers and in particular such as C₁-C₄ alkyl acrylates or
30 methacrylates.

Amphoteric associative polymers according to the invention are
for example described and prepared in patent application WO9844012.

Among the amphoteric associative polymers according to the
invention, the acrylic acid/(meth)acrylamidopropyltrimethylammonium
35 chloride/stearyl methacrylate terpolymers are preferred.

The associative polymers of the known ionic type which can be
used according to the invention are preferably chosen from:

- (1) celluloses modified by groups comprising at least one fatty chain;

there may be mentioned by way of example:

5 - the hydroxyethylcelluloses modified by groups comprising at least one fatty chain such as alkyl, arylalkyl or alkylaryl groups, or mixtures thereof, and in which the alkyl groups are preferably C₈-C₂₂, such as the product NATROSOL PLUS GRADE 330 CS® (C₁₆ alkyls) sold by the company AQUALON, or the product BERMOCOLL EHM 100® sold by the company BEROL NOBEL,

10 - those modified by polyalkylene glycol ether of alkylphenol groups, such as the product AMERCELL POLYMER HM-1500® (polyethylene glycol (15) ether of nonylphenol) sold by the company AMERCHOL.

15 - (2) hydroxypropylguars modified by groups comprising at least one fatty chain such as the product ESAFLOR HM 22® (C₂₂ alkyl chain) sold by the company LAMBERTI, the products RE210-18® (C₁₄ alkyl chain) and RE205-1® (C₂₀ alkyl chain) sold by the company RHONE POULENC.

20 - (3) copolymers of vinylpyrrolidone and of hydrophobic monomers having a fatty chain, of which there may be mentioned by way of example:

- the products ANTARON V216® or GANEX V216® (vinylpyrrolidone/hexadecene copolymer) sold by the company I.S.P.

25 - the products ANTARON V220® or GANEX V220® (vinylpyrrolidone/eicosene copolymer) sold by the company I.S.P.

30 - (4) copolymers of C₁-C₆ alkyl methacrylates or acrylates and of amphiphilic monomers comprising at least one fatty chain such as for example the oxyethylenated stearyl acrylate/methyl acrylate copolymer sold by the company GOLDSCHMIDT under the name ANTIL 208®.

- (5) copolymers of hydrophilic methacrylates or acrylates and of hydrophobic monomers comprising at least one fatty chain such as for example the polyethylene glycol methacrylate/lauryl methacrylate copolymer.

35 - (6) polyether-polyurethanes comprising in their chain both hydrophilic sequences which are most often of a polyoxyethylenated nature and hydrophobic sequences which may be aliphatic chains alone and/or cycloaliphatic and/or aromatic chains.

-(7) polymers containing an aminoplast ether backbone possessing at least one fatty chain, such as the compounds PURE THIX® provided by the company SUD-CHEMIE.

5 Preferably, the polyether-polyurethanes comprise at least two lipophilic hydrocarbon chains, having from 6 to 30 carbon atoms, separated by a hydrophilic sequence, it being possible for the hydrocarbon chains to be pendent chains or chains at the end of a hydrophilic sequence. In particular, it is possible for one or more pendent chains to be envisaged. In addition, the polymer may comprise
10 a hydrocarbon chain at one end or at both ends of a hydrophilic sequence.

The polyether-polyurethanes may be polyblocks, in particular in triblock form. The hydrophobic sequences may be at each end of the chain (for example: triblock copolymer with hydrophilic central
15 sequence) or distributed both at the ends and in the chain (polyblock copolymer for example). These same polymers may also be in the form of graft units or may be star-shaped.

The nonionic polyether-polyurethanes containing a fatty chain may be triblock copolymers whose hydrophilic sequence is a polyoxyethylenated chain comprising from 50 to 1 000 oxyethylenated
20 groups. Nonionic polyether-polyurethanes comprise a urethane bond between the hydrophilic sequences, hence the origin of the name.

By extension, those whose hydrophilic sequences are linked by other chemical bonds to the lipophilic sequences are also included
25 among the nonionic polyether-polyurethanes containing a fatty chain.

By way of examples of nonionic polyether-polyurethanes containing a fatty chain which can be used in the invention, it is also possible to use Rhéolate 205® containing a urea functional group sold by the company RHEOX or the Rhéolates® 208, 204 or 212, as well as
30 Acrysol RM 184®.

There may also be mentioned the product ELFACOS T210® containing a C₁₂₋₁₄ alkyl chain and the product ELFACOS T212® containing a C₁₈ alkyl chain from AKZO.

The product DW 1206B® from RHOM & HAAS containing a
35 C₂₀ alkyl chain and with a urethane bond, sold at 20% dry matter content in water, may also be used.

It is also possible to use solutions or dispersions of these polymers in particular in water or in an aqueous alcoholic medium. By

way of example of such polymers, there may be mentioned Rhéolate® 255, Rhéolate® 278 and Rhéolate® 244 sold by the company RHEOX. It is also possible to use the product DW 1206F and DW 1206J provided by the company ROHM & HAAS.

5 The polyether-polyurethanes which can be used according to the invention are in particular those described in the article by G. Fonnum, J. Bakke and Fk. Hansen - Colloid Polym. Sci 271, 380-389 (1993).

10 Still more particularly it is preferable to use a polyether-polyurethane which can be obtained by polycondensation of at least three compounds comprising (i) at least one polyethylene glycol comprising from 150 to 180 mol of ethylene oxide, (ii) stearyl alcohol or decyl alcohol and (iii) at least one diisocyanate.

15 Such polyether-polyurethanes are sold in particular by the company ROHM & HAAS under the names Aculyn 46® and Aculyn 44® [ACULYN 46® is a polycondensate of polyethylene glycol containing 150 or 180 mol of ethylene oxide, stearyl alcohol and methylenebis(4-cyclohexyl isocyanate) (SMDI), at 15% by weight in a matrix of maltodextrin (4%) and water (81%); ACULYN 44® is a
20 polycondensate of polyethylene glycol containing 150 or 180 mol of ethylene oxide, decyl alcohol and methylenebis(4-cyclohexyl isocyanate) (SMDI), at 35% by weight in a mixture of propylene glycol (39%) and water (26%)].

25 According to a third preferred embodiment, the composition according to the present invention additionally contains at least one surfactant.

 The surfactants which are suitable for carrying out the present invention are in particular the following:

(i) Anionic surfactant(s):

30 By way of example of anionic surfactants which can be used, alone or as mixtures, in the context of the present invention there may be mentioned in particular (nonlimiting list) the salts (in particular alkali metal, especially sodium, salts, ammonium salts, amine salts, amino alcohol salts or magnesium salts) of the following compounds:
35 alkyl sulphates, alkyl ether sulphates, alkylamido ether sulphates, alkylarylpolyether sulphates, monoglyceride sulphates; alkyl sulphonates, alkyl phosphates, alkylamidesulphonates, alkyl aryl sulphonates, α -olefinsulphonates, paraffinsulphonates; (C₆-C₂₄)alkyl

5 sulphosuccinates, (C₆-C₂₄)alkyl ether sulphosuccinates, (C₆-C₂₄)alkylamide sulphosuccinates; (C₆-C₂₄)alkyl sulphoacetates; (C₆-C₂₄)acyl sarcosinates and (C₆-C₂₄)acyl glutamates. It is also possible to use (C₆-C₂₄)alkyl polyglycoside carboxylic esters such as alkyl
10 glucoside citrates, alkyl polyglycoside tartrate and alkyl polyglycoside sulphosuccinates, alkyl sulphosuccinamates; acyl isethionates and N-acyltaurates, the alkyl or acyl radical of all these various compounds preferably comprising from 12 to 20 carbon atoms, and the aryl radical preferably denoting a phenyl or benzyl group. Among the anionic
15 surfactants which can still be used, there may also be mentioned the salts of fatty acids such as the salts of oleic, ricinoleic, palmitic and stearic acids, the acids of copra oil or of hydrogenated copra oil; the acyllactylates whose acyl radical comprises 8 to 20 carbon atoms. It is also possible to use the alkyl D-galactoside uronic acids and their salts, the polyoxyalkylenated (C₆-C₂₄)alkyl ether carboxylic acids, the
20 polyoxyalkylenated (C₆-C₂₄)alkylaryl ether carboxylic acids, the polyoxyalkylenated (C₆-C₂₄)alkyl amido ether carboxylic acids and their salts, in particular those comprising from 2 to 50 alkylene, in particular ethylene, oxide groups, and mixtures thereof.

20 (ii) Nonionic surfactant(s):


The nonionic surfactants themselves are also compounds which are well known per se (in this respect see especially the "Handbook of Surfactants" by M.R. Porter, published by Blackie & Son (Glasgow and London), 1991, pp. 116-178) and, in the context of the present
25 invention, their nature does not assume any critical character. They can thus be chosen especially from (nonlimiting list) alcohols, alpha-diols or polyethoxylated or polypropoxylated alkylphenols which have a fatty chain containing, for example, 8 to 18 carbon atoms, it being possible for the number of ethylene oxide or propylene oxide groups to
30 range especially from 2 to 50. The copolymers of ethylene oxide and propylene oxide and the condensates of ethylene oxide and propylene oxide with fatty alcohols may also be mentioned; the polyethoxylated fatty amides preferably containing from 2 to 30 mol of ethylene oxide, the polyglycerolated fatty amides containing on average 1 to 5
35 glycerol groups and in particular 1.5 to 4; the oxyethylenated fatty acid esters of sorbitan containing from 2 to 30 mol of ethylene oxide; the fatty acid esters of sucrose, the fatty acid esters of polyethylene glycol, alkylpolyglycosides, the N-alkylglucamine derivatives, amine

oxides such as the oxides of (C10-C14)alkylamines or the N-acylaminopropylmorpholine oxides.

(iii) Amphoteric or zwitterionic surfactant(s):


5 The amphoteric or zwitterionic surfactants, the nature of which is not of critical importance in the context of the present invention, may be especially (nonlimiting list) derivatives of aliphatic secondary or tertiary amines in which the aliphatic radical is a linear or branched chain containing 8 to 18 carbon atoms and containing at least one water-solubilizing anionic group (for example carboxylate, sulphonate, sulphate, phosphate or phosphonate); (C₈-C₂₀)alkylbetaines, 10 sulphobetaines, (C₈-C₂₀)alkylamido(C₁-C₆)alkylbetaines or (C₈-C₂₀)alkylamido(C₁-C₆)alkylsulphobetaines may further be mentioned.

15 Among the amine derivatives, there may be mentioned the products sold under the name MIRANOL, as described in patents US-2 528 378 and US-2 781 354 and classified in the CTFA dictionary, 3rd edition, 1982, under the names Amphocarboxyglycinates and Amphocarboxypropionates having the respective structures:



20 in which: R₂ denotes an alkyl radical of an acid R₂-COOH present in hydrolysed copra oil, a heptyl, nonyl or undecyl radical, R₃ denotes a beta-hydroxyethyl group and R₄ a carboxymethyl group;

and



25 in which:

B represents -CH₂CH₂OX', C represents -(CH₂)_z-Y', with z = 1 or 2,

X' denotes the -CH₂CH₂-COOH group or a hydrogen atom

Y' denotes -COOH or the radical -CH₂-CHOH-SO₃H

30 R₂' denotes an alkyl radical of an acid R₉ -COOH present in copra oil or in hydrolysed linseed oil, an alkyl radical, especially C₇, C₉, C₁₁ or C₁₃, a C₁₇ alkyl radical and its iso form or an unsaturated C₁₇ radical.

35 These compounds are classified in the CTFA dictionary, 5th edition, 1993, under the names Disodium Cocoamphodiacetate, Disodium Lauroamphodiacetate, Disodium Caprylamphodiacetate, Disodium Capryloamphodiacetate, Disodium Cocoamphodipropionate, Disodium Lauroamphodipropionate, Disodium Caprylampho-

dipropionate, Disodium Capryloamphodipropionate, Lauro amphodipropionic acid, Cocoamphodipropionic acid.

By way of example, there may be mentioned the cocoamphodiacetate marketed under the trade name MIRANOL® C2M concentrated by the company RHODIA CHIMIE.

(iv) Cationic surfactants:

Among the cationic surfactants, there may be mentioned in particular (nonlimiting list): the salts of optionally polyoxyalkylenated primary, secondary or tertiary amines; quaternary ammonium salts such as tetraalkylammonium, alkylamidoalkyltrialkylammonium, trialkylbenzylammonium, trialkylhydroxyalkylammonium or alkylpyridinium chlorides or bromides; imidazoline derivatives or amine oxides of a cationic nature.

The quantities of surfactants present in the composition according to the invention may vary from 0.01 to 40% and preferably from 0.5 to 30% of the total weight of the composition.

The composition of the present invention may additionally comprise one or more additional oxidation bases which are conventionally used in oxidation dyeing other than the para-phenylenediamines useful for the invention. By way of example, these additional oxidation bases are chosen from phenylenediamines, bisphenylalkylenediamines, ortho-aminophenols, heterocyclic bases other than the heterocyclic para-phenylenediamines useful for the invention and their addition salts.

Among the para-phenylenediamines, there may be mentioned, by way of example, para-phenylenediamine, para-tolylenediamine, 2-chloro-para-phenylenediamine, 2,3-dimethyl-para-phenylenediamine, 2,6-dimethyl-para-phenylenediamine, 2,6-diethyl-para-phenylenediamine, 2,5-dimethyl-para-phenylenediamine, 2- β -hydroxyethyl-para-phenylenediamine, 2-fluoro-para-phenylenediamine, 2-isopropyl-para-phenylenediamine, N-(β -hydroxypropyl)-para-phenylenediamine, 2-hydroxymethyl-para-phenylenediamine, N-(β,γ -dihydroxypropyl)-para-phenylenediamine, N-(4'-aminophenyl)-para-phenylenediamine, N-phenyl-para-phenylenediamine, 2- β -hydroxyethyloxy-para-phenylenediamine, 2- β -acetylaminooethyloxy-para-phenylenediamine, N-(β -methoxyethyl)-para-phenylenediamine, 2-thienyl-para-phenylenediamine, 2- β -

hydroxyethylamino-5-aminotoluene and their addition salts with an acid.

Among the para-phenylenediamines mentioned above, there are particularly preferred para-phenylenediamine, para-tolylenediamine, 2-isopropyl-para-phenylenediamine, 2- β -hydroxyethyl-para-phenylenediamine, 2- β -hydroxyethyloxy-para-phenylenediamine, 2,6-dimethyl-para-phenylenediamine, 2,6-diethyl-para-phenylenediamine, 2,3-dimethyl-para-phenylenediamine, 2-chloro-para-phenylenediamine, 2- β -acetylaminooethyloxy-para-phenylenediamine, and their addition salts with an acid.

Among the bisphenylalkylenediamines, there may be mentioned, by way of example, N,N'-bis(β -hydroxyethyl)-N,N'-bis(4'-aminophenyl)-1,3-diaminopropanol, N,N'-bis(β -hydroxyethyl)-N,N'-bis(4'-aminophenyl)ethylenediamine, N,N'-bis(4-aminophenyl)-tetramethylenediamine, N,N'-bis(β -hydroxyethyl)-N,N'-bis(4-aminophenyl)tetramethylenediamine, N,N'-bis(4-methylaminophenyl)tetramethylenediamine, N,N'-bis(ethyl)-N,N'-bis(4'-amino-3'-methylphenyl)ethylenediamine, 1,8-bis(2,5-diaminophenoxy)-3,6-dioxaoctane, and their addition salts.

Among the ortho-aminophenols, there may be mentioned, by way of example, 2-aminophenol, 2-amino-5-methylphenol, 2-amino-6-methylphenol, 5-acetamido-2-aminophenol, and their addition salts.

Among the heterocyclic bases, there may be mentioned, by way of example, pyridine derivatives, pyrimidine derivatives and pyrazole derivatives.

Among the pyridine derivatives, there may be mentioned the compounds described for example in patents GB 1 026 978 and GB 1 153 196, such as 2,5-diaminopyridine, 2-(4-methoxyphenyl)amino-3-aminopyridine, 3,4-diaminopyridine, and their addition salts.

Other pyridine oxidation bases useful in the present invention are the oxidation bases 3-aminopyrazolo[1,5-a]pyridines or their addition salts which are described, for example, in patent application FR 2801308. By way of example, there may be mentioned pyrazolo[1,5-a]pyridin-3-ylamine; 2-acetylaminopyrazolo[1,5-a]pyridin-3-ylamine; 2-morpholin-4-ylpyrazolo[1,5-a]pyridin-3-ylamine; 3-aminopyrazolo[1,5-a]pyridine-2-carboxylic acid; 2-methoxypyrazolo[1,5-a]pyridin-3-ylamino; (3-aminopyrazolo[1,5-

a]pyridin-7-yl)methanol; 2-(3-aminopyrazolo[1,5-a]pyridin-5-yl)ethanol; 2-(3-aminopyrazolo[1,5-a]pyridin-7-yl)ethanol; (3-aminopyrazolo[1,5-a]pyridin-2-yl)methanol; 3,6-diaminopyrazolo[1,5-a]pyridine; 3,4-diaminopyrazolo[1,5-a]pyridine; pyrazolo[1,5-a]pyridine-3,7-diamine; 7-morpholin-4-ylpyrazolo[1,5-a]pyridin-3-ylamine; pyrazolo[1,5-a]pyridine-3,5-diamine; 5-morpholin-4-ylpyrazolo[1,5-a]pyridin-3-ylamine; 2-[(3-aminopyrazolo[1,5-a]pyridin-5-yl)(2-hydroxyethyl)amino]ethanol; 2-[(3-aminopyrazolo[1,5-a]pyridin-7-yl)(2-hydroxyethyl)amino]ethanol, 3-aminopyrazolo[1,5-a]pyridin-5-ol; 3-aminopyrazolo[1,5-a]pyridin-4-ol, 3-aminopyrazolo[1,5-a]pyridin-6-ol; 3-aminopyrazolo[1,5-a]pyridin-7-ol; and their addition salts.

Among the pyrimidine derivatives, there may be mentioned the compounds described for example in patents DE 2 359 399; JP 88-169 571; JP 05-63124; EP 0 770 375 or patent application WO 96/15765, such as 2,4,5,6-tetraaminopyrimidine, 4-hydroxy-2,5,6-triaminopyrimidine, 2-hydroxy-4,5,6-triaminopyrimidine, 2,4-dihydroxy-5,6-diaminopyrimidine, 2,5,6-triaminopyrimidine, and their addition salts and their tautomeric forms, when a tautomeric equilibrium exists.

Among the pyrazole derivatives, there may be mentioned the compounds described in patents DE 3 843 892, DE 4 133 957 and patent applications WO 94/08969, WO 94/08970, FR-A-2 733 749 and DE 195 43 988 such as 4,5-diamino-1-methylpyrazole, 4,5-diamino-1-(β -hydroxyethyl)pyrazole, 3,4-diaminopyrazole, 4,5-diamino-1-(4'-chlorobenzyl)pyrazole, 4,5-diamino-1,3-dimethylpyrazole, 4,5-diamino-3-methyl-1-phenylpyrazole, 4,5-diamino-1-methyl-3-phenylpyrazole, 4-amino-1,3-dimethyl-5-hydrazinopyrazole, 1-benzyl-4,5-diamino-3-methylpyrazole, 4,5-diamino-3-tert-butyl-1-methylpyrazole, 4,5-diamino-1-tert-butyl-3-methylpyrazole, 4,5-diamino-1-(β -hydroxyethyl)-3-methylpyrazole, 4,5-diamino-1-ethyl-3-methylpyrazole, 4,5-diamino-1-ethyl-3-(4'-methoxyphenyl)pyrazole, 4,5-diamino-1-ethyl-3-hydroxymethylpyrazole, 4,5-diamino-3-hydroxymethyl-1-methylpyrazole, 4,5-diamino-3-hydroxymethyl-1-isopropylpyrazole, 4,5-diamino-3-methyl-1-isopropylpyrazole, 4-amino-5-(2'-aminoethyl)amino-1,3-dimethylpyrazole, 3,4,5-triaminopyrazole, 1-methyl-3,4,5-triaminopyrazole, 3,5-diamino-1-methyl-4-

methylaminopyrazole, 3,5-diamino-4-(β -hydroxyethyl)amino-1-methylpyrazole, and their addition salts.

5 The additional oxidation base(s) present in the composition of the invention are generally present in a quantity of between 0.001 to 20% by weight approximately of the total weight of the dyeing composition, preferably between 0.005 and 6%.

10 The composition according to the invention preferably contains one or more additional couplers conventionally used for dyeing keratinous fibres, other than the benzomorpholine couplers of formula (V') and other than the heterocyclic couplers of formula (V'''). Among these couplers, there may be mentioned in particular meta-phenylenediamines, meta-diphenols, naphthalene couplers, heterocyclic couplers and their addition salts.

15 By way of example, there may be mentioned 1,3-dihydroxybenzene, 1,3-dihydroxy-2-methylbenzene, 4-chloro-1,3-dihydroxybenzene, 2,4-diamino-1-(β -hydroxyethoxy)benzene, 2-amino-4-(β -hydroxyethylamino)-1-methoxybenzene, 1,3-diaminobenzene, 1,3-bis(2,4-diaminophenoxy)propane, 3-ureidoaniline, 3-ureido-1-dimethylaminobenzene, sesamol, 1- β -hydroxyethylamino-3,4-methylenedioxybenzene, α -naphthol, 2-methyl-1-naphthol, 4-hydroxyindole, 4-hydroxy-N-methylindole, 2-amino-3-hydroxypyridine, 3,5-diamino-2,6-dimethoxypyridine, 1-N-(β -hydroxyethyl)amino-3,4-methylenedioxybenzene, 2,6-bis(β -hydroxyethylamino)toluene and their addition salts.

25 In the composition of the present invention, the additional coupler(s) are generally present in a quantity of between 0.001 and 20% by weight approximately of the total weight of the dyeing composition, preferably ranging from 0.005 to 6%.

30 In general, the addition salts of the oxidation bases and couplers which can be used in the context of the invention are in particular chosen from the addition salts with an acid such as the hydrochlorides, hydrobromides, sulphates, citrates, succinates, tartrates, lactates, tosylates, benzenesulphonates, phosphates and acetates and the addition salts with a base such as sodium hydroxide, 35 potassium hydroxide, ammonium hydroxide, amines or alkanolamines.

The dyeing composition in accordance with the invention may additionally contain one or more direct dyes which may be chosen in particular from neutral, acidic or cationic nitro dyes of the benzene

series, neutral, acidic or cation azo direct dyes, neutral, acidic or cationic quinone and in particular anthraquinone direct dyes, azine direct dyes, triarylmethane direct dyes, indoamine direct dyes and natural direct dyes.

5 Among the benzene direct dyes which can be used according to the invention, the following compounds may be mentioned without limitation:

- 1,4-diamino-2-nitrobenzene,
- 1-amino-2 nitro-4- β - hydroxyethylaminobenzene
- 10 - 1-amino-2 nitro-4-bis(β -hydroxyethyl)aminobenzene
- 1,4-bis(β -hydroxyethylamino)-2-nitrobenzene
- 1- β -hydroxyethylamino-2-nitro-4-bis-(β -hydroxyethylamino)benzene
- 1- β -hydroxyethylamino-2-nitro-4-aminobenzene
- 15 - 1- β -hydroxyethylamino-2-nitro-4-(ethyl)(β -hydroxyethyl)aminobenzene
- 1-amino-3-methyl-4- β -hydroxyethylamino-6-nitrobenzene
- 1-amino-2-nitro-4- β -hydroxyethylamino-5-chlorobenzene
- 1,2-diamino-4-nitrobenzene
- 20 - 1-amino-2- β -hydroxyethylamino-5-nitrobenzene
- 1,2-bis(β -hydroxyethylamino)-4-nitrobenzene
- 1-amino-2-tris-(hydroxymethyl)methylamino-5-nitrobenzene
- 1-hydroxy-2-amino-5-nitrobenzene
- 1-hydroxy-2-amino-4-nitrobenzene
- 25 - 1-hydroxy-3-nitro-4-aminobenzene
- 1-hydroxy-2-amino-4,6-dinitrobenzene
- 1- β -hydroxyethyloxy-2- β -hydroxyethylamino-5-nitrobenzene
- 1-methoxy-2- β -hydroxyethylamino-5-nitrobenzene
- 1- β -hydroxyethyloxy-3-methylamino-4-nitrobenzene
- 30 - 1- β , γ -dihydroxypropyloxy-3-methylamino-4-nitrobenzene
- 1- β -hydroxyethylamino-4- β , γ -dihydroxypropyloxy-2-nitrobenzene
- 1- β , γ -dihydroxypropylamino-4-trifluoromethyl-2-nitrobenzene
- 1- β -hydroxyethylamino-4-trifluoromethyl-2-nitrobenzene
- 1- β -hydroxyethylamino-3-methyl-2-nitrobenzene
- 35 - 1- β -aminoethylamino-5-methoxy-2-nitrobenzene
- 1-hydroxy-2-chloro-6-ethylamino-4-nitrobenzene
- 1-hydroxy-2-chloro-6-amino-4-nitrobenzene
- 1-hydroxy-6-bis-(β -hydroxyethyl)amino-3-nitrobenzene

- 1- β -hydroxyethylamino-2-nitrobenzene
- 1-hydroxy-4- β -hydroxyethylamino-3-nitrobenzene.

Among the azo direct dyes which can be used according to the invention, there may be mentioned the cationic azo dyes described in patent applications WO 95/15144, WO 95/01772 and EP-714954 whose content forms an integral part of the invention.

Among these compounds the following dyes may be most particularly mentioned:

- 1,3-dimethyl-2-[[4-(dimethylamino)phenyl]azo]-1H-imidazolium chloride,
- 1,3-dimethyl-2-[(4-aminophenyl)azo]-1H-imidazolium chloride,
- 1-methyl-4-[(methylphenylhydrazono)methyl]pyridinium methyl sulphate.

There may also be mentioned, among the azo direct dyes, the following dyes, which are described in COLOUR INDEX INTERNATIONAL 3rd edition:

- Disperse Red 17
- Acid Yellow 9
- Acid Black 1
- Basic Red 22
- Basic Red 76
- Basic Yellow 57
- Basic Brown 16
- Acid Yellow 36
- Acid Orange 7
- Acid Red 33
- Acid Red 35
- Basic Brown 17
- Acid Yellow 23
- Acid Orange 24
- Disperse Black 9.

There may also be mentioned 1-(4'-aminodiphenylazo)-2-methyl-4bis(β -hydroxyethyl) aminobenzene and 4-hydroxy-3-(2-methoxyphenylazo)-1-naphtalenesulphonic acid.

Among the quinone direct dyes, the following dyes may be mentioned:

- Disperse Red 15

- Solvent Violet 13
- Acid Violet 43
- Disperse Violet 1
- Disperse Violet 4
- 5 -Disperse Blue 1
- Disperse Violet 8
- Disperse Blue 3
- Disperse Red 11
- Acid Blue 62
- 10 -Disperse Blue 7
- Basic Blue 22
- Disperse Violet 15
- Basic Blue 99
- and the following compounds:
- 15 -1-N-methylmorpholiniumpropylamino-4-hydroxyanthraquinone
- 1-aminopropylamino-4-methylaminoanthraquinone
- 1-aminopropylaminoanthraquinone
- 5- β -hydroxyethyl-1,4-diaminoanthraquinone
- 20 -2-aminoethylaminoanthraquinone
- 1,4-bis(β,γ -dihydroxypropylamino)anthraquinone.
- Among the azine dyes, the following compounds may be mentioned:
- Basic Blue 17
- 25 -Basic Red 2.
- Among the triarylmethane dyes which can be used according to the invention, the following compounds may be mentioned:
- Basic Green 1
- Acid blue 9
- 30 -Basic Violet 3
- Basic Violet 14
- Basic Blue 7
- Acid Violet 49
- Basic Blue 26
- 35 -Acid Blue 7
- Among the indoamine dyes which can be used according to the invention, the following compounds may be mentioned:

- 2- β -hydroxyethylamino-5-[bis(β -4'-hydroxyethyl)amino]anilino-1,4-benzoquinone
- 2- β -hydroxyethylamino-5-(2'-methoxy-4'-amino)anilino-1,4-benzoquinone
- 5 - 3-N(2'-chloro-4'-hydroxy)phenylacetyl amino-6-methoxy-1,4-benzoquinoneimine
- 3-N(3'-chloro-4'-methylamino)phenylureido-6-methyl-1,4-benzoquinoneimine
- 10 - 3-[4'-N-(ethylcarbamylmethyl)amino]phenylureido-6-methyl-1,4-benzoquinoneimine

Among the natural direct dyes which can be used according to the invention, there may be mentioned lawsone, juglone, alizarin, purpurin, carminic acid, kermesic acid, purpurogallin, protocatechaldehyde, indigo, isatin, curcumin, spinulosin and apigenidin. It is also possible to use extracts or decoctions containing these natural dyes and in particular henna-based poultices or extracts.

The direct dye(s) preferably represent from 0.001 to 20% by weight approximately of the total weight of the ready-to-use composition and still more preferably from 0.005 to 10% by weight approximately.

The composition according to the invention may also contain at least one hydroxylated solvent, such as in particular ethanol, propylene glycol, glycerol, polyol monoethers, benzyl alcohol.

It may also contain a nonhydroxylated solvent.

The hydroxylated solvents and the nonhydroxylated solvents are preferably present in proportions preferably between 1 and 40% by weight approximately relative to the total weight of the dyeing composition, and still more preferably between 5 and 30% by weight approximately.

The dyeing composition in accordance with the invention may also contain various adjuvants conventionally used in hair dyeing compositions, such as antioxidants, penetrating agents, sequestering agents, perfumes, buffers, dispersing agents, conditioning agents such as for example modified or unmodified, volatile or nonvolatile silicones, film-forming agents, ceramides, preservatives and opacifying agents.

The above adjuvants are generally present in a quantity, for each of them, of between 0.01 and 20% by weight relative to the weight of the composition.

Of course, persons skilled in the art will be careful to choose this or these optional additional compounds such that the advantageous properties intrinsically attached to the oxidation dyeing composition in accordance with the invention are not, or not substantially, impaired by the addition(s) envisaged.

The pH of the dyeing composition in accordance with the invention is generally between 3 and 12 approximately, and preferably between 5 and 11 approximately. It may be adjusted to the desired value by means of acidifying or alkalinizing agents customarily used in dyeing keratinous fibres or with the aid of conventional buffering systems.

Among the acidifying agents, there may be mentioned, by way of example, inorganic or organic acids such as hydrochloric acid, orthophosphoric acid, sulphuric acid, carboxylic acids such as acetic acid, tartaric acid, citric acid, lactic acid and sulphonic acids.

Among the alkalinizing agents, there may be mentioned, by way of example, ammonium hydroxide, alkali metal carbonates, alkanolamines such as mono-, di- and triethanolamines and the derivatives thereof, sodium or potassium hydroxides and the compounds of the following formula (XXIII):



(XXIII)

in which W is a propylene residue optionally substituted with a hydroxyl group or a C₁-C₄ alkyl radical; R_a, R_b, R_c and R_d, which may be identical or different, represent a hydrogen atom, a C₁-C₄ alkyl radical or a C₁-C₄ hydroxyalkyl radical.

The dyeing composition according to the invention may be provided in various forms, such as in the form of liquids, creams or gels, or in any other appropriate form for dyeing keratinous fibres, and in particular human hair.

The method of the present invention is a method in which the composition according to the present invention, as defined above, is

5 applied to the fibres, and the colour is developed using an oxidizing agent. The colour may be developed at acidic, neutral or alkaline pH and the oxidizing agent may be added to the composition of the invention just at the time of use or it can be used from an oxidizing composition containing it, applied simultaneously or sequentially with the composition of the invention.

10 According to a particular embodiment, the composition according to the present invention is mixed, preferably at the time of use, with a composition containing, in a medium appropriate for dyeing, at least one oxidizing agent, this oxidizing agent being present in a sufficient quantity to develop a colour. The mixture obtained is then applied to the keratinous fibres. After an exposure time of 3 to 50 minutes approximately, preferably 5 to 30 minutes approximately, the keratinous fibres are rinsed, washed with shampoo, rinsed again and then dried.

15 The oxidizing agents conventionally used for the oxidation dyeing of keratinous fibres are, for example, hydrogen peroxide, urea peroxide, alkali metal bromates, persalts such as perborates and persulphates, peracids and the oxidase enzymes, among which there may be mentioned peroxidases, oxidoreductases with 2 electrons such as uricases and oxygenases with 4 electrons such as laccases. Hydrogen peroxide is particularly preferred.

20 The oxidizing composition may also contain various adjuvants conventionally used in hair dyeing compositions and as defined above.

25 The pH of the oxidizing composition containing the oxidizing agent is such that, after mixing with the dyeing composition, the pH of the resulting composition applied to keratinous fibres preferably varies between 3 and 12 approximately, and still more preferably between 5 and 11. It may be adjusted to the desired value by means of acidifying or alkalinizing agents customarily used for dyeing keratinous fibres and as defined above.

30 The ready-to-use composition which is finally applied to the keratinous fibres may be provided in various forms, such as in the form of liquids, creams or gels, or in any other form appropriate for dyeing keratinous fibres, and in particular human hair.

35 The subject of the invention is finally a multicompartment device or dyeing "kit" in which a first compartment contains the dyeing composition defined above and a second compartment contains

an oxidizing composition. This device may be equipped with means which make it possible to deliver the desired mixture to the hair, such as the devices described in patent FR-2 586 913 in the name of the applicant.

Using this device, it is possible to dye keratinous fibres using a method which comprises mixing a dyeing composition in accordance with the invention with an oxidizing agent as defined above, and applying the mixture obtained to the keratinous fibres for a time sufficient to develop the desired colour.

DETAILED DESCRIPTION

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventors to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

Example 1 :

Dyeing composition: (expressed in grams)

Oleyl alcohol	6
Oleic acid	3
Polyglycerolenated oleyl alcohol containing 2 moles of glycerol	6
Polyglycerolenated oleyl alcohol containing 6 moles of glycerol	6
Diethylaminopropyl laurylaminosuccinamate, sodium salt	3
Oxyethylenated oleyl amine containing 2 moles of ethylene oxide	7
Monoethanolamide of alkyl ether carboxylic acid containing 2 moles of ethylene oxide	10
Ammonium acetate	20
Propylene glycol	20

Reducing agents, antioxidants	0.915
Sequestrants	1
Para-aminophenol	0.1
[1-(4-Aminophenyl)pyrrolidin-3-yl]trimethylammonium chloride	0.3
N,N-(2-Hydroxyethyl)-p-phenylenediamine sulphate	0.3
6-Hydroxybenzomorpholine	0.45
Perfume	qs
Ammonium hydroxide (containing 20.5% ammonia)	10.2
Demineralized water	qs 100

At the time of use, this composition is mixed weight for weight with an oxidizing milk containing 6% hydrogen peroxide. The mixture obtained is applied for 30 minutes to grey hair which is 90% white. A flat dark blond coloration is obtained on this hair after rinsing, shampooing and drying.

Example 2 :

The same result is obtained if the molar equivalent of the cationic tertiary base in this example is replaced by 3-[1-(4-amino-phenyl)pyrrolidin-3-yl]-1-methyl-3H-imidazol-1-ium chloride.

Example 3 :

Dyeing composition: (expressed in grams)

Oleyl alcohol	6
Oleic acid	3
Polyglycerolenated oleyl alcohol containing 2 moles of glycerol	6
Polyglycerolated oleyl alcohol containing 6 moles of glycerol	6
Diethylaminopropyl laurylamino succinamate, sodium salt	3
Oxyethylenated oleyl amine containing 2 moles of ethylene oxide	7
Alkyl ether carboxylic acid monoethanolamide containing 2 moles of ethylene oxide	10

Ammonium acetate	20
Propylene glycol	20
Reducing agents, antioxidants	0.915
Sequestrants	1
N,N-(2-Hydroxyethyl)-p-phenylenediamine sulphate	0.3
[1-(4-Aminophenyl)pyrrolidin-3-yl]trimethyl-ammonium chloride	0.3
6-Hydroxybenzomorpholine	0.15
6-Hydroxyindole	0.15
Perfume	qs
Ammonium hydroxide (containing 20.5% ammonia)	10.2
Demineralized water	qs 100

At the time of use, this composition is mixed weight for weight with an oxidizing milk containing 6% hydrogen peroxide. The mixture obtained is applied for 30 minutes to grey hair which is 90% white. A dark blond coloration is obtained on this hair after rinsing, shampooing and drying.